

Review on recent results of J/ψ production at STAR

Jitka Mrázková^{1,2,*} for the STAR Collaboration

¹*Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague*

²*Nuclear Physics Institute, Czech Academy of Sciences*

**Address correspondence to: jitka.mrazkova@fjfi.cvut.cz*

Abstract

Studying the production of J/ψ (bound state of charm and anti-charm quark) in proton-proton collisions gives an opportunity to test quantum chromodynamics (QCD) calculations, as the production of J/ψ involves both perturbative and non-perturbative processes. However, theoretical calculations are still unable to fully and simultaneously explain experimental results, such as polarization and p_T spectra measured in different kinematic regimes and colliding energies. More studies are needed to investigate J/ψ production mechanism. In heavy-ion collisions, charmonia can be used to study the properties of the medium as they are expected to dissociate in the medium when the Debye radius, inversely proportional to the medium temperature, becomes smaller than their size. Other competing effects, such as recombination, have also been found to modify the observed J/ψ yield in heavy-ion collisions. We review recent measurements of the J/ψ production in proton-proton and heavy-ion collisions at various collision energies measured with the STAR experiment at RHIC. The data are compared with recent model calculations on charmonia production.

1 Introduction

Understanding the production mechanisms of J/ψ mesons is crucial for testing QCD calculations in proton-proton ($p+p$) collisions and probing the properties of the strongly interacting medium created in heavy-ion collisions. J/ψ suppression provides evidence of QGP formation, where color screening prevents the binding of charm quarks, depending on the energy density and temperature of the medium [1]. Describing charmonium production in a medium is challenging due to the competing effects of recombination and dissociation processes. Systematic studies of various collision systems and energies may help to disentangle the charmonium production mechanism. Accordingly, the following proceedings provide an overview of recent results on the J/ψ production from the STAR experiment.

2 Experimental setup

All of the analyses presented here were obtained from data collected using the STAR (Solenoidal Tracking at RHIC) detector [2]. Some of the key detectors utilized in J/ψ -related analyses at mid-rapidity include the Time Projection Chamber (TPC) [3], which provides charged particle tracking, momentum determination, and energy loss measurements for particle identification. The Time-of-Flight (TOF) detector [4] is used for complementary particle identification. The Barrel Electromagnetic Calorimeter (BEMC) [5] operates as for particle detection based on deposited energy, provides high- p_T triggering, and features fine granularity in $(\eta, \varphi) = (0.05, 0.05)$ with full azimuthal coverage: $0 \leq \varphi < 2\pi$.

3 Results

Firstly, the results focused on suppression of the J/ψ production, reconstructed via the di-electron decay channel, in heavy-ion collisions across various collision systems and energies will be presented. This will be followed by a discussion of the analyses related to the J/ψ production in proton-proton collisions, including studies of multiplicity dependence and the J/ψ production within jets.

3.1 Inclusive J/ψ R_{AA}

To study the influence of the medium on the J/ψ production, one can use the nuclear modification factor R_{AA} defined as:

$$R_{AA} = \frac{\sigma_{\text{inel}}}{\langle N_{\text{coll}} \rangle} \frac{d^2 N_{AA}/dydp_T}{d^2 \sigma_{pp}/dydp_T}, \quad (1)$$

where $\langle N_{\text{coll}} \rangle$ corresponds to a scaling factor representing an average number of binary nucleon-nucleon collisions based on the Glauber model predictions [6]. N_{AA} is the average number of J/ψ produced in one minimum-bias A+A collisions and σ_{pp} is the cross section in non-single diffractive (NSD) p+p collisions, as a function of transverse momentum p_T and rapidity y .

Fig. 1a shows the dependence of the J/ψ R_{AA} on collision energy in the range of $\sqrt{s_{NN}} = 14.6\text{--}200$ GeV in Au+Au collisions [7, 8], and includes a comparison with SPS [9], ALICE data [10, 11] and model predictions [12, 13]. Energy dependence can be qualitatively described by the transport model [5,6], with primordial production being dominant at RHIC energies and regeneration at the LHC [7,8], as illustrated by the model curves in the figure. No significant energy dependence of J/ψ R_{AA} is observed in central collisions within uncertainties of up to 200 GeV.

Another measurement, the measurement of J/ψ R_{AA} as a function of centrality in Au+Au collisions at different collision energies, is presented in Fig. 1b, where the J/ψ R_{AA} indicates a slightly decreasing trend, hence suggesting a stronger suppression in central collisions compared to the peripheral collisions. No significant energy dependence is observed in a given $\langle N_{part} \rangle$.




Figure 1: (a) J/ψ R_{AA} as a function of collision energy with data from the STAR experiment $\sqrt{s_{NN}} = 14.6\text{--}200$ GeV in Au+Au collisions [7, 8], compared with SPS and ALICE measurements [9, 10, 11] and model predictions [12, 13]. (b) Measurement of the J/ψ R_{AA} as a function of centrality in Au+Au collisions for various collision energies at the STAR experiment.

3.2 $\psi(2S)$ over J/ψ Double Ratio

The first observation of sequential charmonium suppression in heavy-ion collisions at STAR is shown in the Fig. 2 and quantified by the $\psi(2S)$ over J/ψ double ratio:

$$\frac{[(\text{Bd}\sigma_{\psi(2S)}) / (\text{Bd}\sigma_{J/\psi})]_{AA}}{[(\text{Bd}\sigma_{\psi(2S)}) / (\text{Bd}\sigma_{J/\psi})]_{pp,pd}}, \quad (2)$$

which is described by the production cross sections of the excited charmonium state $\psi(2S)$ and of the ground state J/ψ (i.e. $\sigma_{\psi(2S)}$ and $\sigma_{J/\psi}$, respectively) and their branching ratios B , comparing heavy-ion (A+A) collisions to proton-proton ($p+p$) or proton-deuteron ($p+d$) collisions.

It is observed that $\psi(2S)$ is overall more suppressed than J/ψ , reflecting its weaker binding energy and greater sensitivity to the temperature of the QGP. Suppression grows with centrality, driven by higher energy density and QGP lifetime in central collisions. The data are compared with an average $p+p$ reference from NA51, ISR and PHENIX [14, 15, 16]. The double ratio appears to be smaller in the isobar system than in the $p+A$ system (see the right panel of the Fig. 2). Other measurements for different collision systems and energies are further shown for comparison.




Figure 2: $\psi(2S)$ over J/ψ double ratio as a function of N_{part} for various A+A collision systems (left), and the results for $p+A$ systems (right panel). The data are compared with an average $p+p$ reference from NA51, ISR and PHENIX [14, 15, 16].

3.3 J/ψ Production vs Multiplicity in $p + p$

76

In high multiplicity $p + p$ collisions, multiple parton interactions (MPI) and string percolation effects are expected to play a significant role in modifying the bulk production. These mechanisms can enhance the initial parton density and increase the probability of interactions, potentially affecting the yield of the produced J/ψ mesons.

77

78

79

80

The measured data are compared to previous results at $\sqrt{s} = 200$ GeV [17] in Fig. 3. The J/ψ normalized yields to their event average at $\sqrt{s} = 510$ GeV are consistent with those at $\sqrt{s} = 200$ GeV, with a wider multiplicity range achieved at $\sqrt{s} = 510$ GeV. There is an indication of a splitting between the results obtained at RHIC and LHC energies [18, 19]. The difference between LHC and RHIC results is still under investigation.

81

82

83

84

85




Figure 3: The measurement of the J/ψ normalized yields (relative to their event-average) vs normalized charged particle multiplicity in $p + p$ collisions for different collision energies.

3.4 J/ψ Production in Jets in p + p collisions

The study of J/ψ production in jets provides additional discriminative power for production mechanisms. The fraction z of charged-particle jet transverse momentum carried by J/ψ is defined as the ratio of the J/ψ transverse momentum to the transverse momentum of the jet:

$$z(J/\psi) = \frac{p_T^{J/\psi}}{p_T^{\text{jet}}}. \quad (3)$$

The measured z distribution for inclusive J/ψ in jets in $p + p$ collisions at $\sqrt{s} = 500$ GeV normalized by the J/ψ cross-section [20], is compared to model prediction (PYTHIA 8) and shown in Fig. 4. The results show discrepancy with the model predictions. The z distribution remains relatively flat, while PYTHIA predicts a steep rise toward $z = 1$, where most the jet momentum is carried by the J/ψ.




Figure 4: The normalized z distributions for inclusive J/ψ mesons produced within a jet in $p + p$ collisions at $\sqrt{s} = 500$ GeV compared to prediction from PYTHIA 8. The data are normalized by the J/ψ cross-section at the same collision energy [20].

4 Summary and Outlook

97

In these proceedings, we have shown the recent measurements of charmonium production have been shown in A + A collisions, including the study of J/ψ R_{AA} and the charmonium sequential suppression using the $\psi(2S)$ over J/ψ double ratio. The J/ψ production dependence on charged-particle multiplicity at $\sqrt{s} = 510$ GeV and its production in jets in $p + p$ collisions at $\sqrt{s} = 500$ GeV was presented.

98

99

100

101

102

Further charmonia measurements have not been covered in this paper, such as the study of azimuthal anisotropy and polarization. Studies of J/ψ polarization in jets in $p + p$ collisions are ongoing to provide deeper insights into the J/ψ production mechanism. The high luminosity $p + p$ and Au+Au data at 200 GeV from 2023–2025 [21] will enable more precise measurements of J/ψ elliptic anisotropy and $\psi(2S)$ production, as suggested in Fig. 5.

103

104

105

106

107




Figure 5: Projection of $\psi(2S)$ to J/ψ ratios vs centrality for the planned 2023–2025 data-taking period at the STAR experiment [21].

References

108

- [1] S. Digal, P. Petreczky, and H. Satz. *Sequential Quarkonium Suppression*. *Physics Letters B*, 514:57–62, **2001**.
- [2] STAR Collaboration. *STAR Detector Overview*. *Nucl. Instrum. Meth. A*, 499:624–632, **2003**.

109

110

111

112

- [3] Anderson et al. *The STAR Time Projection Chamber: A Unique Tool for Studying High Multiplicity Events at RHIC. Nucl. Instrum. Meth. A*, 499:659–678, **2003**. 113
114
- [4] Ackermann et al. *Extensive particle identification with TPC and TOF at the STAR experiment. Nucl. Instrum. Meth. A*, 558:419–429, **2005**. 115
116
- [5] Beddo et al. *The STAR Barrel Electromagnetic Calorimeter. Nucl. Instrum. Meth. A*, 499:725–739, **2003**. 117
118
- [6] Miller et al. *Glauber Modeling in High-Energy Nuclear Collisions. Ann. Rev. Nucl. Part. Sci.*, 57:205–243, **2007**. 119
120
- [7] STAR Collaboration. *Energy dependence of J/ψ production in Au+Au collisions at $\sqrt{s_{NN}} = 39, 62.4$ and 200 GeV. Phys. Lett. B*, 771:13, **2017**. 121
122
- [8] STAR Collaboration. *Measurement of inclusive J/ψ suppression in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV through the dimuon channel at STAR. Phys. Lett. B*, 797:134917, **2019**. 123
124
125
- [9] NA50 Collaboration. *Evidence for Deconfinement of Quarks and Gluons from the J/ψ Suppression Pattern Measured in Pb-Pb Collisions at the CERN SPS. Phys. Lett. B*, 477:28, **2000**. 126
127
128
- [10] ALICE Collaboration. *Centrality, rapidity, and transverse momentum dependence of J/ψ suppression in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. Phys. Lett. B*, 734:314, **2014**. 129
130
- [11] ALICE Collaboration. *Quarkonium measurements in nucleus-nucleus collisions with ALICE. Phys. Lett. B*, 1005:121769, **2021**. 131
132
- [12] X. Zhao and R. Rapp. *Charmonium in Medium: From Correlators to Experiment. Phys. Rev. C*, 82:064905, **2010**. 133
134
- [13] L. Kluberg. *20 years of J/ψ suppression at the CERN SPS. Eur. Phys. J. C*, 43:145, **2005**. 135
136
- [14] PHENIX Collaboration. *Ground and excited charmonium state production in $p + p$ collisions at $\sqrt{s} = 200$ GeV. Phys. Rev. D*, 85:092004, **2012**. 137
138
- [15] NA51 Collaboration. *J/ψ , ψ' and Drell-Yan production in pp and pd interactions at 450 GeV/c. Phys. Lett. B*, 438:35, **1998**. 139
140
- [16] CERN ISR. *Electron pair production at the CERN ISR. Nucl. Phys. B*, 142:29, **1978**. 141
- [17] STAR Collaboration. *J/ψ production cross section and its dependence on charged-particle multiplicity in $p + p$ collisions at $\sqrt{s} = 200$ GeV. Phys. Lett. B*, 786:87, **2018**. 142
143

- [18] ALICE Collaboration. *Multiplicity dependence of J/ψ production at midrapidity in pp collisions at $\sqrt{s} = 13 \text{ TeV}$. *Phys. Lett. B*, 810:135758, **2020**.*
- [19] ALICE Collaboration. *J/ψ Production as a Function of Charged Particle Multiplicity in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$. *Phys. Lett. B*, 712:165, **2012**.*
- [20] STAR Collaboration. *Measurements of the transverse-momentum-dependent cross sections of production at mid-rapidity in proton+proton collisions at $\sqrt{s} = 510$ and 500 GeV with the STAR detector. *Phys. Rev. D*, 100:052009, **2019**.*
- [21] STAR Collaboration. *STAR Beam Use Request Runs 24 - 25*. STAR BUR Runs 24 - 25. Accessed 2023.