

Recent studies on heavy-flavor femtoscopy in heavy-ion collisions by STAR

Priyanka Roy Chowdhury (for the STAR Collaboration)

Warsaw University of Technology, Poland

Heavy quarks are produced in hard partonic scatterings at the very early stage of heavy-ion collisions and they experience the whole evolution of the Quark-Gluon Plasma medium. Femtoscopic correlations, i.e. two-particle correlations at low relative momentum, are sensitive to the final-state interactions as well as to the extent of the region from which the correlated particles are emitted. A study of correlations between heavy-flavor mesons and identified charged hadrons could shed light on their interactions in the hadronic phase.

STAR has performed the first measurement of femtoscopic correlation between D^0 -hadron pairs at mid-rapidity in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV. D^0 mesons are reconstructed via the $K^{\mp} - \pi^{\pm}$ decay channel using topological criteria enabled by the Heavy Flavor Tracker with excellent track pointing resolution. We will present the femtoscopic correlation functions between $D^0/\overline{D^0}$ - π^{\pm} , $D^0/\overline{D^0}$ - K^{\pm} and $D^0/\overline{D^0}$ - p^{\pm} pairs for $D^0/\overline{D^0}$ with transverse momentum above 1 GeV/c in the 0 – 80% centrality range. STAR results will be compared with existing theory predictions and its physics implications will also be discussed.