

1 Azimuthal Transverse Single-Spin Asymmetries of Identified
2 Hadrons Within Jets from Polarized pp Collisions at $\sqrt{s} =$
3 200 GeV

4 Ting Lin, for the STAR Collaboration
5 Shandong University

6 May 26, 2021

7 **Abstract**

8 The Collins effect involves the convolution of the quark transversity in the proton with the
9 spin-dependent Collins fragmentation function, leading to azimuthal modulations of identified
10 charged hadron yields relative to the jet axis. Recently, a detailed calculation using the soft-
11 collinear effective theory found that the Collins effect in pp collisions involves a mixture of
12 collinear and transverse momentum dependent (TMD) factorization. It provides a direct
13 probe of the Collins fragmentation function and enables testing of its evolution, universality
14 and factorization breaking in the transverse momentum dependent formalism. In 2018, STAR
15 published the first measurements of Collins asymmetries for charged pions in jets in polarized
16 pp collisions at $\sqrt{s} = 500$ GeV based on data taken during 2011. The results probe Q^2 scales
17 one to two orders of magnitude larger than similar measurements in semi-inclusive deep-
18 inelastic scattering (SIDIS) and are consistent with predictions based on global analyses
19 of e^+e^- and SIDIS data. In 2012 and 2015, STAR collected ~ 14 pb $^{-1}$ and ~ 48 pb $^{-1}$ of
20 transversely polarized pp data at $\sqrt{s} = 200$ GeV, respectively. These datasets provide the
21 most precise measurement of the Collins effect in pp collisions to date, especially at the quark
22 momentum fractions $0.1 \leq x \leq 0.4$. Preliminary results for Collins asymmetries of identified
23 pions, kaons, and protons in jets at $\sqrt{s} = 200$ GeV and comparisons to theory predictions
will be presented.