

 Brookhaven[™]
National Laboratory

 APS Topical Group on
Hadronic Physics
GHP

11th Workshop of the APS Topical Group
on Hadronic Physics

Recent results on vector meson photoproduction and interference effects in Ultra Peripheral Collisions at STAR

Ashik Ikbal Sheikh
(For the STAR Collaboration)

 KENT STATE
UNIVERSITY

Kent State University

March 14 - 16, 2025, Anaheim, USA


Supported in part by



U.S. DEPARTMENT OF
ENERGY

Office of
Science

Heavy ions miss each other: Ultra-Peripheral Collisions (UPCs)




Collisions where nuclei do NOT collide

No hadronic collisions happen

Ions interact through photon-ion
and photon-photon collisions

=> Called Ultra-peripheral
collisions (UPCs)

The strongest EM-fields in UPCs



● In UPCs,

$$E_{max} = 10^{18} \text{ V/m}, B_{max} \sim 10^{14} - 10^{18} \text{ T}$$

=> Strongest EM-field in the universe, but transient

● EM-field treated in terms of quasi-real photons



$$E_{\gamma,max} \sim \gamma \hbar c / R;$$

$$E_{\gamma,max} \sim 30 \text{ GeV (RHIC@Au+Au 200 GeV)}$$

$$E_{\gamma,max} \sim 80 \text{ GeV (LHC@Pb+Pb 2.76 TeV)}$$

=> EM-fields are quantized as photons in UPCs


Photon-gluon scattering: Vector meson (VM) production via photon-nuclear interactions



Photoproduction of Vector Mesons (VM) in UPC


UPC VM: Powerful probe of parton densities inside nuclei

Satre simulation of parton density fluctuations, Fig: A. Kumar




$\lambda_{coherent}$

Low p_T



$\lambda_{incoherent}$

High p_T





● Probes parton density & fluctuations inside nuclei—
constraints for A+A collisions initial state

● Modification of parton densities in heavy nuclei



=> VM helps to probe parton density inside nuclei before EIC era

UPC events with STAR detector





- Neutron(s) detected in ZDCs
- ZDC signals show peak structure for neutrons
- No activity in both BBCs => Diffractive events (η -gap)

=> Method to trigger UPC events



=> Coherent and incoherent contributions can be disentangled via the combined fit of mass and p_T




Rapidity dependence J/ψ production cross-section

STAR, *Phys Rev Lett* 133 (2024) 5, 052301

- Measured for coherent and incoherent contributions for different neutron emission in ZDCs
- Systematic uncertainties in incoherent to coherent cross-section ratio are largely cancelled
- Sensitive to the nuclear structure and deformation

=> Important to constrain theoretical models related to nuclear geometry





J/ψ Nuclear suppression factors

STAR, *Phys Rev Lett* 133 (2024) 5, 052301

- Coherent cross-section suppressed by ~30% w.r.t free nucleon
- The incoherent supp. is ratio b/w incoh x-sec with HERA (H1) free proton data
- Incoherent photoproduction has been suppressed by ~65% (at $W_{\gamma^*N} = 25$ GeV) w.r.t free proton H1 data
- Stronger incoherent suppressions than model predictions — Even does not directly support the CGC with subnucleonic fluctuations

=> Provides constraints to the parton density and baseline for future measurements in EIC






VM spin interference: A novel quantum phenomenon for high resolution gluon imaging


Polarized Photons from colliding nuclei

STAR, Phys. Rev. Lett. 127 (2021) 52302




Transverse view of Lorentz contracted nuclei

=> Photons in UPC are linearly polarized — polarization is roughly along impact parameter




Experimental access to photon polarization demonstrated by STAR, measuring the Breit-Wheeler process, $\gamma\gamma \rightarrow e^+e^-$

UPC vector meson spin and decay daughters are correlated



=> The $\cos(2\phi)$ modulation in VM momentum distribution w.r.t photon polarization direction


Measuring the modulation over a large no. of events




Photon polarization correlated with Impact parameter \rightarrow random from one event to the next

\Rightarrow Event average washes out the $\cos(2\phi)$ modulation w.r.t photon polarization direction

Photon source ambiguity





PATH - 1



PATH - 2

\Rightarrow Two independent paths of VM production
 \rightarrow The paths are indistinguishable

Interference makes the modulation observable in experiment





Double Slit Experiment

Best analogy: Double slit experiment in Optics

=> Two indistinguishable paths may interfere and make the $\cos(2\phi)$ modulation observable

Photon source ambiguity: Interference among amplitudes of two possible paths

Observation of interference for $\rho^0 \rightarrow \pi^+\pi^-$ at STAR



SCIENCE ADVANCES | RESEARCH ARTICLE

PHYSICS STAR, Sci. Adv. 9, eabq 3903 (2023)

Tomography of ultrarelativistic nuclei with polarized photon-gluon collisions

STAR Collaboration


A linearly polarized photon can be quantized from the Lorentz-boosted electromagnetic field of a nucleus traveling at ultrarelativistic speed. When two relativistic heavy nuclei pass one another at a distance of a few nuclear radii, the photon from one nucleus may interact through a virtual quark-antiquark pair with gluons from the other nucleus, forming a short-lived vector meson (e.g., ρ^0). In this experiment, the polarization was used in diffractive photoproduction to observe a unique spin interference pattern in the angular distribution of $\rho^0 \rightarrow \pi^+\pi^-$ decays. The observed interference is a result of an overlap of two wave functions at a distance an order of magnitude larger than the ρ^0 travel distance within its lifetime. The strong-interaction nuclear radii were extracted from these diffractive interactions and found to be 6.53 ± 0.06 fm (^{197}Au) and 7.29 ± 0.08 fm (^{238}U), larger than the nuclear charge radii. The observable is demonstrated to be sensitive to the nuclear geometry and quantum interference of nonidentical particles.

Copyright © 2023 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).


Observed the interference for coherent ρ^0 photoproduction in UPCs

Measured in 3 different collision systems: Au+Au, U+U, p+Au \rightarrow Sensitive to nuclear shape/size

The p_T dependence of interference for $\rho^0 \rightarrow \pi^+\pi^-$ at STAR



STAR: Signal $\pi^+\pi^-$ pairs



SCIENCE ADVANCES | RESEARCH ARTICLE

PHYSICS STAR, Sci. Adv. 9, eabq 3903 (2023)

Tomography of ultrarelativistic nuclei with polarized photon-gluon collisions

STAR Collaboration

A linearly polarized photon can be quantized from the Lorentz-boosted electromagnetic field of a nucleus traveling at ultrarelativistic speed. When two relativistic heavy nuclei pass one another at a distance of a few nuclear radii, the photon from one nucleus may interact through a virtual quark-antiquark pair with gluons from the other nucleus, forming a short-lived vector meson (e.g., ρ^0). In this experiment, the polarization was used in diffractive photoproduction to observe a unique spin interference pattern in the angular distribution of $\rho^0 \rightarrow \pi^+\pi^-$ decays. The observed interference is a result of an overlap of two wave functions at a distance an order of magnitude larger than the ρ^0 travel distance within its lifetime. The strong-interaction nuclear radii were extracted from these diffractive interactions and found to be 6.53 ± 0.06 fm (^{197}Au) and 7.29 ± 0.08 fm (^{238}U), larger than the nuclear charge radii. The observable is demonstrated to be sensitive to the nuclear geometry and quantum interference of nonidentical particles.

Copyright © 2023 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).


Clear p_T dependence of interference observed

Interference gets weak at higher p_T —
Incoherent processes take over


Radius measurement with interference for $\rho^0 \rightarrow \pi^+\pi^-$ at STAR

STAR, Sci. Adv. 9, eabq 3903 (2023)


C STAR: Au+Au $\sqrt{s_{NN}}=200$ GeV



D STAR: Au+Au $\sqrt{s_{NN}}=200$ GeV



A STAR: Photonuclear $\rho^0 \rightarrow \pi^+\pi^-$




Impact of spin interference on $|t|$ distribution studied in different ϕ bins

Improved measurement of mass radii using spin interference effect

$R(\text{Au}) = 6.53 \pm 0.06$ fm; $R(\text{U}) = 7.29 \pm 0.08$ fm

Spin interference with $J/\psi \rightarrow e^+e^-$




$J/\psi \rightarrow e^+e^-$


Boson Fermions

Measured sign of the interference tells us whether the interference occurring in daughter or parent level

Interference of quantum particles \rightarrow Spin interference




Mass: 0.7 GeV/c²
Lifetime: 1.3 fm/c




Mass: 3.1 GeV/c²
Lifetime: 2160 fm/c

J/ψ heavier than ρ^0 and J/ψ has longer lifetime




Probes finer structure and captures high quality images of the gluon distributions

Measured spin interference with $J/\psi \rightarrow e^+e^-$



Observable for J/ψ spin interference



Interference signal fitted with: $1 + a_2 \cos(2\phi) \Rightarrow a_2$ is the measure of the modulation



Measured $\cos(2\phi)$ for spin interference of J/ψ s

Observed spin interference for $J/\psi \rightarrow e^+e^-$

The p_T -dependent interference of J/ ψ





Diff+Int predictions : Mäntysaari et al. Phys.Rev.C 109 (2024) 2, 024908

- Interference signal for J/ ψ shows p_T dependence
- Positive modulation for ρ and negative for J/ ψ ($a_2 \sim -12\%$ with 3σ for $p_T < 100$ MeV)
- Diffractive+interference calculations cannot describe the data well

\Rightarrow Useful for gluon tomography within the nucleus

Decay anisotropy of photo-produced J/ψ in heavy ion peripheral collisions



=> Significant modulation ($\sim 39\%$) w.r.t reaction plane

=> Probes photon polarization and the initial collision geometry


Summary and take home

- STAR Measured the coherent and incoherent J/ψ production in Au+Au UPCs
- STAR observed the spin interference of the photoproduced ρ^0 and J/ψ
- Measured interference signal has p_T dependence
- Measured the photon induced J/ψ polarization w.r.t reaction plane in peripheral collisions
- Measurements are sensitive to nuclear geometry and useful to constrain the theoretical models
- RHIC, LHC and future EIC experiments can provide further insights into these

Thank You!

Backup

STAR detector



- Main central barrel detectors for UPC measurements: TPC, TOF, BEMC
- Forward detectors: BBC or EPD, ZDC

Incoherent J/ψ production cross-section vs p_T^2

STAR, *Phys Rev Lett* 133 (2024) 5, 052301

○ Incoherent production compared with H1 data with free proton


○ Strong nuclear suppression ($\sim 49\%$) seen
(Mäntysaari et. al, *Phys. Rev. Lett.* 117 (2016) 5, 052301)

○ Models found H1 data supports sub-nucleonic fluctuations




(Mäntysaari et. al, *Phys. Rev. D* 106 (2022) 7, 074019)

○ STAR data shows the bound nucleon has similar shape as the free proton — similar sub-nucleonic fluctuations in heavy nuclei

=> Strong nuclear suppression and sub-nucleonic fluctuations in Au nucleus



Corrections for interference signal



◎ The $\gamma + \gamma \rightarrow e^+ + e^-$ has also the J/Ψ interference like pattern due to detector effect

◎ Correct for the 2γ process with : $a_2 = f \times a_2^{bkg} + (1 - f) \times a_2^{sig}$, with $f = \frac{N_{bkg}}{N_{sig} + N_{bkg}}$

◎ Considered the Bremsstrahlung process and $J/\psi \rightarrow e^+ + e^- + \gamma$, using the STARLight+Geant simulations

=> Background correction is done to extract true modulation signal