



Electric charge and strangeness dependent splitting of the rapidity-odd directed flow in Au+Au collisions

Ashik Ikbal Sheikh (for the STAR Collaboration)

Kent State University





Directed flow (v_1) and splitting (Δv_1)

- ▶ First harmonic coefficient of Fourier decomposition of particle azimuthal distribution, v_1 - Directed Flow

$$E \frac{d^3 N}{dp^3} = \frac{d^2 N}{2\pi p_T dp_T dy} \left(1 + 2 \sum_{n=1}^{+\infty} v_n \cos[n(\phi - \Psi_{RP})] \right)$$


where $v_n = \langle \cos[n(\phi - \Psi_{RP})] \rangle$

- ▶ Probe early stage of the collisions - strong EM-Field




- ▶ What drives the splitting - Initial EM-Field or QCD-driven effect?
- ▶ Measure splitting with charge (Δq) and strangeness (ΔS)

EM-Field driven splitting (Δv_1) - Faraday and Hall effect?



- ▶ Beam direction: \hat{z} and Impact parameter: $\hat{x} \Rightarrow$ Reaction Plane: xz
- ▶ Colliding nuclei produce B-field, \perp to RP (approx) \Rightarrow B along \hat{y}
- ▶ Time varying \vec{B} induces \vec{E} field \Rightarrow Faraday effect
- ▶ Medium expands longitudinally ($\vec{u} \perp \vec{B}$) - Lorentz force pushes +ve and -ve charged particles in opposite directions \Rightarrow Hall effect

EM-Field driven splitting (Δv_1)?




U. Gursoy et al., Phys. Rev. C 89, 054905 (2014)

- ▶ Faraday and Hall are competing effects - Net effect affects v_1
- ▶ v_1 for +ve particles shown (when Faraday > Hall)

Multi-strange and the splitting (Δv_1)

- ▶ Enhanced strange quarks production and identity retains during hadronization => multiply multi-strange baryons (Ξ and Ω)
- ▶ Low scattering cross section and early thermal freeze-out - good probe of early stage of the collisions
- ▶ Multi-strange v_1 might be important for strangeness related splitting


Towards measurements: STAR detector at BES-II



Journal of Physics: Conference Series 742 (2016) 012022

- ▶ TPC+TOF for PID: TPC measures $-dE/dx$ of tracks ($|\eta| < 1$, $0 < \phi < 2\pi$) and TOF measures time of flight ($|\eta| < 0.9$)
- ▶ EPD ($2.1 < |\eta| < 5.1$) or ZDC ($|\eta| > 6.3$) for event plane reconstruction
- ▶ Data sets (analyzed):
Au+Au at $\sqrt{s_{NN}} = 27$ GeV (year-2018) and $\sqrt{s_{NN}} = 200$ GeV (year-2016)

Splitting (Δv_1): Choice of particles?



(1) Measurements with heavy flavors?

- ▶ Measurements of HFs are challenging
- ▶ Less abundantly produced - suffer large uncertainties
- ▶ Absence of HFT in STAR BES-II and low production rate - HF measurements are difficult

(2) Measurements with light hadrons?



- ▶ Light hadrons produced in abundance - precise measurements
- ▶ Δv_1 measurements come with drawbacks:
 - (a) Most of the (anti)-particles contain transported quarks (u and d)
 - (b) Transported quarks have different v_1 than the produced \Rightarrow Δv_1 becomes difficult to interpret
- ▶ Avoiding transported quarks \Rightarrow Splitting can be measured with light hadrons

Splitting (Δv_1): Our Approach

- ▶ Use only produced particles, K^- , \bar{p} , $\bar{\Lambda}$, ϕ , Ξ^+ , Ω^- and $\bar{\Omega}^+$
- ▶ Based on Quark coalescence
- ▶ Coalescence-inspired sum rule: $v_1(\text{Hadron}) = \sum v_1^i(q_i)$
- ▶ A new way to test coalescence sum rule (same $y - p_T/n_q$ phase space, with $n_q \rightarrow$ no. of constituent quarks):

$$v_1[K^-(\bar{u}s)] + v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] = v_1[\bar{p}(\bar{u}\bar{u}\bar{d})] + v_1[\phi(s\bar{s})] \quad (1)$$

$$v_1[K^-(\bar{u}s)] + v_1[\Xi^+(\bar{d}\bar{s}\bar{s})] = v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] + v_1[\phi(s\bar{s})] \quad (2)$$





Splitting (Δv_1): Our Approach

- ▶ New idea to show the coalescence sum rule holds (with identical quarks):

$$v_1[K^-(\bar{u}s)] + v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] = v_1[\bar{p}(\bar{u}\bar{u}\bar{d})] + v_1[\phi(s\bar{s})] \quad (1)$$

$$v_1[K^-(\bar{u}s)] + v_1[\Xi^+(\bar{d}\bar{s}\bar{s})] = v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] + v_1[\phi(s\bar{s})] \quad (2)$$



- ▶ With produced particles, K^- , \bar{p} , $\bar{\Lambda}$, ϕ , Ξ^+ , Ω^- and $\bar{\Omega}^+$ and make combinations - having same quark mass but different Δq and ΔS





Rearranging the Δv_1 in Δq and ΔS

Particles: $K^-(\bar{u}s)$, $\bar{p}(\bar{u}\bar{u}\bar{d})$, $\bar{\Lambda}(\bar{u}\bar{d}\bar{s})$, $\phi(s\bar{s})$, $\Xi^+(\bar{d}\bar{s}\bar{s})$, $\Omega^-(sss)$, $\bar{\Omega}^+(\bar{s}\bar{s}\bar{s})$

Index	Quark Mass	Charge	Strangeness	Expression
1	$\Delta m = 0$	$\Delta q = 0$	$\Delta S = 0$	$[\bar{p}(\bar{u}\bar{u}\bar{d}) + \phi(s\bar{s})] - [K^-(\bar{u}s) + \bar{\Lambda}(\bar{u}\bar{d}\bar{s})]$
2	$\Delta m \approx 0$	$\Delta q = \frac{2}{3}$	$\Delta S = 1$	$[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [\frac{1}{2}\phi(s\bar{s}) + \frac{2}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$
3	$\Delta m \approx 0$	$\Delta q = 1$	$\Delta S = 2$	$[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [\frac{1}{3}\Omega^-(sss) + \frac{2}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$
4	$\Delta m \approx 0$	$\Delta q = \frac{4}{3}$	$\Delta S = 2$	$[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [K^-(\bar{u}s) + \frac{1}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$
5	$\Delta m \approx 0$	$\Delta q = \frac{4}{3}$	$\Delta S = 2$	$[\Xi^+(\bar{d}\bar{s}\bar{s})] - [\phi(s\bar{s}) + \frac{1}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$
6	$\Delta m = 0$	$\Delta q = 2$	$\Delta S = 6$	$[\bar{\Omega}^+(\bar{s}\bar{s}\bar{s})] - [\Omega^-(sss)]$
7	$\Delta m \approx 0$	$\Delta q = \frac{7}{3}$	$\Delta S = 4$	$[\Xi^+(\bar{d}\bar{s}\bar{s})] - [K^-(\bar{u}s) + \frac{1}{3}\Omega^-(sss)]$


- ▶ Combinations have same $\Delta m(\approx 0)$ different Δq and ΔS - 7 combinations
- ▶ Degenerate combinations (Indices 4 and 5) - Good cross check
- ▶ Measure splitting with Δq and ΔS

v_1 vs y : Ξ and Ω Baryons




- ▶ $\Xi(\Lambda\pi)$ and $\Omega(\Lambda K)$ baryons reconstructed by KF-Particle
- ▶ $v_1(y)$ for multi-strange hadrons - First measurement!
- ▶ Large v_1 for Ω baryons - the statistical uncertainties are large

Splitting (Δv_1) at 3 different Δq and ΔS (27 GeV)




- ▶ Δv_1 for same mass, different charge and strangeness
- ▶ Δv_1 increases at larger y for $\Delta q \neq 0$
- ▶ Δv_1 also increases with p_T/n_q when $\Delta q \neq 0$
- ▶ AMPT (Phys. Rev. C 100, 054903 (2019)) has opposite trend for $\Delta q \neq 0$ - No EM-Field is implemented in AMPT

Δv_1 -slope - splitting: hints of QED and/or QCD effect



- ▶ Δv_1 -slope ($d\Delta v_1/dy$) increases with increasing Δq and ΔS at 27 GeV
- ▶ Δq and ΔS are correlated (see Table at page-9)
- ▶ For 27 GeV, slope = 0.002905 ± 0.000481 (with Δq); $> 5\sigma$ effect
- ▶ For 200 GeV, slope = 0.001159 ± 0.00038 (with Δq); $> 2.5\sigma$ effect
- ▶ $d\Delta v_1/dy$ -slope is less for 200 GeV than 27 GeV

Δv_1 -slope - splitting: Model comparison



- ▶ AMPT can not explain the data (Phys. Rev. C 100, 054903 (2019))
- ▶ PHSD(+EM-Field) can describe the data within the uncertainties

Summary

- ▶ First measurements of v_1 of multi-strange baryons - Ξ and Ω
- ▶ Measured charge (Δq) and strangeness (ΔS) dependent splitting, Δv_1 , at BES-II
- ▶ Δv_1 -slope ($d\Delta v_1/dy$) increases as Δq and ΔS increase at 27 GeV
- ▶ PHSD+EM-Field calculations can describe data within uncertainties - Hints of EM-Field effect in the splitting
- ▶ Net strangeness is also an important key factor for Δv_1 -slope

THANK YOU