## Central Exclusive Production with the STAR detector at RHIC

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

Leszek Adamczyk AGH University of Krakow for the STAR Collaboration

## Abstract

We present results on the Central Exclusive Production of charged hadron pairs  $h^+h^-(h = \pi, K, p)$ obtained in the STAR experiment at RHIC in proton-proton collisions at center-of-mass energies of  $\sqrt{s} = 200$  GeV and 510 GeV. All final-state particles of the process  $pp \rightarrow p' + h^+h^- + p'$  were reconstructed, including forward-scattered protons detected in the Roman Pot system. As a result, the Double Pomeron Exchange (DPE) events were selected and the non-exclusive backgrounds were efficiently rejected. The DPE process can be regarded as a spin and parity filter, i.e., the  $h^+h^-$  system must have even spin and positive parity. It is gluon-rich process a favorable place to look for scalar glueball states.

At  $\sqrt{s} = 200$  GeV differential fiducial cross sections were measured as functions of observables related to the central hadronic final state and to the forward-scattered protons. The measured cross sections were compared to phenomenological predictions based on the DPE model. Structures observed in the mass spectra of  $\pi^+\pi^-$  and  $K^+K^-$  pairs were found consistent with the DPE model, while angular distributions of pions suggested a dominant spin-0 contribution to  $\pi^+\pi^-$  production. We also present preliminary results on the measurement of the same physics process at higher  $\sqrt{s} = 510$  GeV.