

# Strangeness and electric charge dependent splitting of the rapidity-odd directed flow in Au+Au collisions

Ashik Ikbal Sheikh (for the STAR Collaboration)

Kent State University

2021 Fall Meeting of the APS Division of Nuclear Physics  
(DNP 2021)

Massachusetts Institute of Technology (MIT)  
Virtual Meeting, October 11–14, 2021





## Directed flow ( $v_1$ ) and splitting ( $\Delta v_1$ )

- ▶ First harmonic coefficient of Fourier decomposition of particle azimuthal distribution,  $v_1$  - Directed Flow

$$E \frac{d^3 N}{dp^3} = \frac{d^2 N}{2\pi p_T dp_T dy} \left( 1 + 2 \sum_{n=1}^{+\infty} v_n \cos[n(\phi - \Psi_{RP})] \right)$$


where  $v_n = \langle \cos[n(\phi - \Psi_{RP})] \rangle$

- ▶ Probe early stage of the collisions - strong EM-Field




- ▶ What drives the splitting - Initial EM-Field or QCD-driven effect?
- ▶ Measure splitting with charge ( $\Delta q$ ) and strangeness ( $\Delta S$ )

# EM-Field driven splitting ( $\Delta v_1$ ) - Faraday and Hall effect?



- ▶ Beam direction:  $\hat{z}$  and Impact parameter:  $\hat{x} \Rightarrow$  Reaction Plane:  $xz$
- ▶ Colliding nuclei produce B-field,  $\perp$  to RP (approx)  $\Rightarrow$  B along  $\hat{y}$
- ▶ Time varying  $\vec{B}$  induces  $\vec{E}$  field  $\Rightarrow$  Faraday effect
- ▶ Medium expands longitudinally ( $\vec{v} \perp \vec{B}$ ) - Lorentz force pushes +ve and -ve charged particles in opposite directions  $\Rightarrow$  Hall effect

# EM-Field driven splitting ( $\Delta v_1$ )?




U. Gursoy et al., Phys. Rev. C 89, 054905 (2014)

- ▶ Faraday and Hall are competing effects - Net effect affects  $v_1$
- ▶  $v_1$  for +ve particles shown (when Faraday > Hall)

## Multi-strange and the splitting ( $\Delta v_1$ )

- ▶ Enhanced strange quarks production and identity retains during hadronization => multiply multi-strange baryons ( $\Xi$  and  $\Omega$ )
- ▶ Low scattering cross section and early thermal freeze-out - good probe of early stage of the collisions
- ▶ Multi-strange  $v_1$  might be important for strangeness related splitting


# Towards measurements: STAR detector at BES-II



Journal of Physics: Conference Series 742 (2016) 012022

- ▶ TPC+TOF for PID: TPC measures  $-dE/dx$  of tracks ( $|\eta| < 1$ ,  $0 < \phi < 2\pi$ ) and TOF measures time of flight ( $|\eta| < 0.9$ )
- ▶ EPD ( $2.1 < |\eta| < 5.1$ ) or ZDC ( $|\eta| > 6.3$ ) for event plane reconstruction
- ▶ Data sets (analyzed):  
Au+Au at  $\sqrt{s_{NN}} = 27$  GeV (year-2018) and  $\sqrt{s_{NN}} = 200$  GeV (year-2016)

# Splitting ( $\Delta v_1$ ): Choice of particles?



## (1) Measurements with heavy flavors?

- ▶ Measurements of HFs are challenging
- ▶ Less abundantly produced - suffer large uncertainties
- ▶ Absence of HFT in STAR BES-II and low production rate - HF measurements are difficult

## (2) Measurements with light hadrons?


- ▶ Light hadrons produced in abundance - precise measurements
- ▶  $\Delta v_1$  measurements come with drawbacks:
  - (a) Most of the (anti-)particles contain transported quarks (u and d)
  - (b) Transported quarks have different  $v_1$  than the produced  $\Rightarrow$   $\Delta v_1$  becomes difficult to interpret
- ▶ Avoiding transported quarks  $\Rightarrow$  Splitting can be measured with light hadrons

## Splitting ( $\Delta v_1$ ): Our Approach

- ▶ Use only produced particles,  $K^-$ ,  $\bar{p}$ ,  $\bar{\Lambda}$ ,  $\phi$ ,  $\Xi^+$ ,  $\Omega^-$  and  $\bar{\Omega}^+$
- ▶ Based on Quark coalescence
- ▶ Coalescence-inspired sum rule:  $v_1(\text{Hadron}) = \sum v_1^i(q_i)$
- ▶ A new way to test coalescence sum rule (same  $y - p_T/n_q$  phase space, with  $n_q \rightarrow$  no. of constituent quarks):

$$v_1[K^-(\bar{u}s)] + v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] = v_1[\bar{p}(\bar{u}\bar{u}\bar{d})] + v_1[\phi(s\bar{s})] \quad (1)$$

$$v_1[K^-(\bar{u}s)] + v_1[\Xi^+(\bar{d}\bar{s}\bar{s})] = v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] + v_1[\phi(s\bar{s})] \quad (2)$$



$$\Omega^-(sss) \quad \bar{\Omega}^+(\bar{s}\bar{s}\bar{s})$$

# Splitting ( $\Delta v_1$ ): Our Approach



- ▶ New idea to show the coalescence sum rule holds (with identical quarks):

$$v_1[K^-(\bar{u}s)] + v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] = v_1[\bar{p}(\bar{u}\bar{u}\bar{d})] + v_1[\phi(s\bar{s})] \quad (1)$$

$$v_1[K^-(\bar{u}s)] + v_1[\Xi^-(\bar{d}\bar{s}\bar{s})] = v_1[\bar{\Lambda}(\bar{u}\bar{s}\bar{d})] + v_1[\phi(s\bar{s})] \quad (2)$$

(1)

(2)



- ▶ With produced particles,  $K^-$ ,  $\bar{p}$ ,  $\bar{\Lambda}$ ,  $\phi$ ,  $\Xi^-$ ,  $\Omega^-$  and  $\bar{\Omega}^+$  and make combinations - having same quark mass but different  $\Delta q$  and  $\Delta S$







## Rearranging the $\Delta v_1$ in $\Delta q$ and $\Delta S$

Particles:  $K^-(\bar{u}s)$ ,  $\bar{p}(\bar{u}\bar{u}\bar{d})$ ,  $\bar{\Lambda}(\bar{u}\bar{d}\bar{s})$ ,  $\phi(s\bar{s})$ ,  $\Xi^+(\bar{d}\bar{s}\bar{s})$ ,  $\Omega^-(sss)$ ,  $\bar{\Omega}^+(\bar{s}\bar{s}\bar{s})$

| Index | Quark Mass           | Charge                   | Strangeness    | Expression   |
|-------|----------------------|--------------------------|----------------|--|
| 1     | $\Delta m = 0$       | $\Delta q = 0$           | $\Delta S = 0$ | $[\bar{p}(\bar{u}\bar{u}\bar{d}) + \phi(s\bar{s})] - [K^-(\bar{u}s) + \bar{\Lambda}(\bar{u}\bar{d}\bar{s})]$       |
| 2     | $\Delta m \approx 0$ | $\Delta q = \frac{2}{3}$ | $\Delta S = 1$ | $[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [\frac{1}{2}\phi(s\bar{s}) + \frac{2}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$ |
| 3     | $\Delta m \approx 0$ | $\Delta q = 1$           | $\Delta S = 2$ | $[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [\frac{1}{3}\Omega^-(sss) + \frac{2}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$  |
| 4     | $\Delta m \approx 0$ | $\Delta q = \frac{4}{3}$ | $\Delta S = 2$ | $[\bar{\Lambda}(\bar{u}\bar{d}\bar{s})] - [K^-(\bar{u}s) + \frac{1}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$             |
| 5     | $\Delta m \approx 0$ | $\Delta q = \frac{4}{3}$ | $\Delta S = 2$ | $[\Xi^+(\bar{d}\bar{s}\bar{s})] - [\phi(s\bar{s}) + \frac{1}{3}\bar{p}(\bar{u}\bar{u}\bar{d})]$                    |
| 6     | $\Delta m = 0$       | $\Delta q = 2$           | $\Delta S = 6$ | $[\bar{\Omega}^+(\bar{s}\bar{s}\bar{s})] - [\Omega^-(sss)]$  |
| 7     | $\Delta m \approx 0$ | $\Delta q = \frac{7}{3}$ | $\Delta S = 4$ | $[\Xi^+(\bar{d}\bar{s}\bar{s})] - [K^-(\bar{u}s) + \frac{1}{3}\Omega^-(sss)]$                                      |


- ▶ Combinations have same  $\Delta m(\approx 0)$  different  $\Delta q$  and  $\Delta S$  - 7 independent combinations
- ▶ Degenerate combinations (Indices 4 and 5) - Good cross check
- ▶ Measure splitting with  $\Delta q$  and  $\Delta S$

# $v_1$ vs $y$ : $\Xi$ and $\Omega$ Baryons




- ▶  $\Xi(\Lambda\pi)$  and  $\Omega(\Lambda K)$  baryons reconstructed by KF-Particle
- ▶  $v_1$  ( $y$ ) for multi-strange hadrons - First measurement!
- ▶ Large  $v_1$  for  $\Omega$  baryons - the statistical uncertainties are large

# Splitting ( $\Delta v_1$ ) at 3 different $\Delta q$ and $\Delta S$ (27 GeV)




- ▶  $\Delta v_1$  for same mass, different charge and strangeness
- ▶  $\Delta v_1$  increases at larger  $y$  for  $\Delta q \neq 0$
- ▶  $\Delta v_1$  also increases with  $p_T/n_q$  when  $\Delta q \neq 0$
- ▶ AMPT ([Phys. Rev. C 100, 054903 \(2019\)](#)) has opposite trend for  $\Delta q \neq 0$  - No EM-Field is implemented in AMPT

# $\Delta v_1$ -slope - splitting: hints of QED and/or QCD effect



- ▶  $\Delta v_1$ -slope ( $d\Delta v_1/dy$ ) increases with increasing  $\Delta q$  and  $\Delta S$  at 27 GeV
- ▶  $\Delta q$  and  $\Delta S$  are correlated (see Table at page-9)
- ▶ For 27 GeV, slope =  $0.002905 \pm 0.000481$  (with  $\Delta q$ );  $> 5\sigma$  effect
- ▶ For 200 GeV, slope =  $0.001159 \pm 0.00038$  (with  $\Delta q$ );  $> 2.5\sigma$  effect
- ▶  $d\Delta v_1/dy$ -slope is less for 200 GeV than 27 GeV

## $\Delta v_1$ -slope - splitting: Model comparison



- ▶ AMPT can not explain the data (Phys. Rev. C 100, 054903 (2019))
- ▶ PHSD(+EM-Field) can describe the data within the uncertainties

## Summary

- ▶ First measurements of  $v_1$  of multi-strange baryons -  $\Xi$  and  $\Omega$
- ▶ Measured charge ( $\Delta q$ ) and strangeness ( $\Delta S$ ) dependent splitting,  $\Delta v_1$ , at BES-II
- ▶  $\Delta v_1$ -slope ( $d\Delta v_1/dy$ ) increases as  $\Delta q$  and  $\Delta S$  increase at 27 GeV
- ▶ PHSD+EM-Field calculations can describe data within uncertainties - Hints of EM-Field effect in the splitting
- ▶ Net strangeness is also an important key factor for  $\Delta v_1$ -slope

THANK YOU