

Measurement of Mid-rapidity Inclusive Jet Cross Section in pp Collisions at $\sqrt{s} = 200$ GeV

Dmitry Kalinkin
For the STAR Collaboration

¹Indiana University – Bloomington

²Brookhaven National Laboratory

2020 Fall Meeting of the APS Division of Nuclear Physics
30 October 2020



Supported in part by




U.S. DEPARTMENT OF
ENERGY

Office of
Science





Proton Structure in Hard Interactions



Jet production in high energy collisions of hadrons can be described in terms of following ingredients:


- Initial state of hadrons
- Hard collision of partons
- Parton Shower
- Underlying Event (UE)
- Hadronization

$$d\sigma_{pp \rightarrow jet+X}(Q^2) = \sum_{a,b} \int \underbrace{f_a(x_1, Q^2) f_b(x_2, Q^2)}_{\text{proton structure}} \underbrace{d\hat{\sigma}_{a+b \rightarrow jet+X}(x_1, x_2, Q^2)}_{\text{hard process+PS+Had.}} dx_1 dx_2$$




Original plot from NNPDF 3.1 Catalog of plots: α_s variations at NNLO

Jet Measurements using STAR Detector





Original illustration by Tai Sakuma

- **TPC:** Interaction vertex and charged particle tracks
- **BEMC** and **EEMC:** Photon energy measurement
- **Trigger condition** on deposited EM energy sum in 1×1 patches in $\eta - \phi$
- East and west **Zero Degree Calorimeter:** Absolute luminosity monitoring




Gluon Polarization using Jets at STAR


- Measurements using a similar collinear factorization framework $A_{LL} \sim \Delta f_a \otimes \Delta f_b \otimes \Delta \hat{\sigma}$ to determine Δg the helicity distribution of gluons inside the proton
- Detector effects are not unfolded but corrected by adjusting p_T (or M_{jj}) and A_{LL} of independent points
- Run 12 mid-rapidity inclusive and di-jet A_{LL} results recently published [PRD 100 (2019) 052005]
- Run 13 and Run 15 publications coming soon



E. R. Nocera at Transversity 2017,




PRD 100 (2019) 114027




PRD 100 (2019) 052005

Published Jet Cross Sections from STAR



Phys. Rev. Lett. 97 (2006) 252001

- An inclusive jet cross section
- Mid-point cone algorithm
- Not corrected for UE or hadronization
- Bin-by-bin detector effects correction
- Limited acceptance




Phys Rev D 95 (2017) 071103

- A di-jet cross section
- anti- k_T algorithm
- Detector effects unfolded
- No data-driven UE correction


Jets at Three Levels

Parton jets




- Made of partons outgoing from the hard interaction
- Definition flexible depending on theoretical needs (e.g. fit using pQCD)

Particle jets



- Made of stable particles (at STAR the π^0 treated as stable)
- Universal – Free from all detector effects
- Includes effects of
 - QCD radiation
 - Hadronization
 - UE (unless subtracted)

Detector jets




- Made of tracks and discrete calorimeter towers
- Experiment specific

©2009 Tai Sakuma


Detector Smearing Unfolding

Particle jets




Simulation with Pythia 6 and
GEANT3

Unfolding of p_T spectra by
inverting the detector
response matrix:



Detector jets




©2009 T. Geurts

- Matrix inversion gives the exact result for the maximum likelihood estimator
- Statistical fluctuations are regularized by choosing sufficiently large bin sizes
- **Need to estimate uncertainty due to the choice of prior (in this case, Pythia)**

Underlying Event Correction

- Two off-axis cone regions defined as $(\varphi - \varphi_{\text{jet}} \pm \pi/2)^2 + (\eta - \eta_{\text{jet}})^2 \leq R_{\text{UE}}^2$ with $R_{\text{UE}} = 0.5$





- For each jet calculate a jet area A and a p_T -density of constituents ρ_{UE}
- Correction implemented via a jet p_T shift:

$$\text{jet } p_T \rightarrow \text{jet } p_T - A \cdot \rho_{\text{UE}}$$

- Applied to data before unfolding and to simulation in definition of the detector response

Charged UE measured at STAR:




Phys Rev D 101 (2020) 052004

(Regions defined as $|\varphi - \varphi_{\text{jet}} \pm \pi/2| < \pi/6$)


Uncertainties for Incl. Jet Cross Section at $\sqrt{s} = 200$ GeV

Leading systematic uncertainties are due to:

- Jet Energy Scale
 - BEMC response – **the dominant uncertainty!**
 - Calibration
 - Response to charged hadrons
 - TPC
 - Tracking efficiency
 - Resolution
- Luminosity measurement
- Hadronization correction
- UE correction
- Simulation sample statistics
- Choice of event generator used to simulate the detector response (estimated to be insignificant)



Inclusive Jet Cross Section at $\sqrt{s} = 510$ GeV, Particle Level





Different triggers:

- **JP0**: $E \geq 5.4$ GeV
- **JP1**: $E \geq 7.3$ GeV
- **JP2**: $E \geq 14.4$ GeV

Measured in two η -ranges:

- $0 < |\eta| < 0.5$
- $0.5 < |\eta| < 0.9$



Conclusions

- Jet measurements at STAR are extended to the unpolarized case
- Inclusive jet measurements at RHIC will allow to better constrain high- x behaviour of the gluon PDF
- ...and serve as a normalization for other possible measurements like measurement of hadron fragmentation inside jets

$$\left(\frac{d^2\sigma}{dp_{T;\text{jet}} dz_h} \right) / \left(\frac{d\sigma}{dp_{T;\text{jet}}} \right)$$

- Measurements at two values of \sqrt{s} , at 200 GeV and 510 GeV, provide insights into energy dependence of the Underlying Event