

The splitting of rapidity-odd directed flow of produced particles in Au+Au and isobar collisions at STAR

Ashik Ikbal Sheikh (for the STAR collaboration)

Department of Physics, Kent State University, Kent, OH 44242, USA

1 Rapidity-odd directed flow (v_1) of hadrons in relativistic heavy-ion collisions can provide insights into the ultra-
2 strong electromagnetic (EM) field [1-2], the constituent quark content of hadrons [3], and the role of transported
3 quarks [4]. The v_1 for identified charged hadrons with unprecedented precision is reported in Au+Au and isobar
4 collisions at $\sqrt{s_{NN}} = 200$ GeV.

5 The coalescence sum rule is examined with various combinations of hadrons where all constituent quarks are
6 produced. For such combinations a systematic violation of the sum rule is observed with increasing difference in the
7 electric charge and the strangeness content of the associated constituent quarks. By comparing with the Parton-
8 Hadron-String Dynamics model that includes an EM field, the results suggest that the constituent quark sum rule
9 could be violated in the presence of a strong EM field that drives the v_1 of produced quarks and anti-quarks to opposite
10 directions. The splitting of v_1 slope with rapidity ($\Delta(dv_1/dy)$) between positively and negatively charged hadrons
11 (π , K , p) is also studied with large statistics. A clear transition of $\Delta(dv_1/dy)$ from positive in central collisions to
12 negative in peripheral collisions is observed for protons and kaons. With the effects of both transported quarks and
13 the EM field considered, it is found that the significant negative values in peripheral events can only be explained by
14 the presence of an EM field in the collisions.

15
16
17 [1] U. Gursoy, et al., *Phy. Rev. C* **98**, 055201 (2018).

18 [2] U. Gursoy, et al., *Phy. Rev. C* **89**, 054905 (2014).

19 [3] A. I. Sheikh, et al., *Phy. Rev. C* **105**, 014912 (2022).

20 [4] Y. Guo, et al. *Phys. Rev. C* **86**, 044901 (2012).