

# Dilepton production in p+p, Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and U+U collisions at $\sqrt{s_{NN}} = 193$ GeV

---

DANIEL BRANDENBURG  
RICE UNIVERSITY  
FOR THE STAR COLLABORATION



## Part I

- Very low  $p_T$   $e^+e^-$  invariant mass spectra in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV and U+U collisions at  $\sqrt{s_{NN}} = 193$  GeV

**See Shuai Yang's Poster : Poster ID 340**

## Part II

- First look at  $\mu^+\mu^-$  invariant mass spectra with the Muon Telescope Detector at STAR in p+p collisions at  $\sqrt{s} = 200$  GeV and peripheral Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV

# Motivation: Low $p_T$ $J/\psi$ Enhancement



→ STAR and ALICE have observed significant excess production of  $J/\psi$  in **peripheral** A+A collisions at **low  $p_T$**  ( $p_T < 300$  MeV/c)


→ Cannot be described by hadronic production modified by medium / cold nuclear matter effects

→ May be due to coherent photoproduction?

→ Observed in collisions with impact parameter  $b < 2 R$

**Motivation to measure  $e^+e^-$  pair production over a wider invariant mass range**

$J/\psi$   $R_{AA}$  in Au+Au and U+U Collisions via  $J/\psi \rightarrow e^+e^-$



See Wangmei Zha's poster  
Poster ID 258


## Time Projection Chamber (TPC)

- Charged particle tracking
- Momentum measurement
- Particle identification via ionization energy loss ( $dE/dx$ )

## Time-Of-Flight (TOF) Detector


- Used in  $e^+e^-$  analysis to reject slow hadrons
- Enables clean electron identification up to  $p_T \sim 3 \text{ GeV}/c$

$0 < p_T < 0.15 \text{ GeV}/c$




- Significant excess in 60-80% central Au+Au and U+U collisions for  $M_{ee} < 4 \text{ GeV}/c^2$
- Excess becomes less significant in more central collisions

# STAR $e^+e^-$ in Au+Au & U+U STAR



- Excess is consistent in Au+Au and U+U
- Excess cannot be explained by hadronic contributions modified by medium
- Can this excess be explained by coherent photoproduction?






→ Good agreement with hadronic cocktail contributions for  $p_T > 0.2 \text{ GeV/c}$

→ Excess is found entirely below  $p_T < 0.2 \text{ GeV/c}$

# STAR $e^+e^-$ in Au+Au & U+U STAR




- Low  $p_T$  shape very similar to that from coherent photoproduction in Ultra Peripheral Collisions
- Can this excess in  $e^+e^-$  be explained by coherent photoproduction?



→ What is coherent photoproduction?

Photoproduction in Ultra Peripheral Collisions (UPC)




C. Bertulani, S. Klein, J. Nystrand Ann. Rev. Nucl. Part. Sci.55:271(2005)

Coherent:

- Emitted photon/pomeron interacts with the nucleus as a whole
- Strong coupling ( $Z\alpha_{EM} \sim 0.6$ ) results in large cross sections
- Photon wavelength  $\lambda = h/p > R_A$
- $p_T < h/R_A \sim 30 \text{ MeV}/c$  for heavy ions

# Photon Interactions in A+A Collisions

- We can gain insight into the production mechanism through the  $t = p_T^2$  distribution due to interference effects



S. Klein and J. Nystrand, PRL **84** 2330 (2000)

Coherent photon-nucleus interaction:

- Photon can be emitted from  $A_1$  or  $A_2$ , ie. two indistinguishable processes
- For a vector meson, the amplitudes from each process have opposite signs
- Results in strong **destructive interference** at very low  $p_T$
- Experimentally distinguish by looking at  $dN/dp_T^2$  distribution at very low  $p_T$

# Coherent Photoproduction in UPCs STAR

→ What does this look like in the UPC case?


→ Coherent production characterized by **large slope parameter** for small  $|t|$

$$\text{SLOPE} = 426 \pm 2 \text{ (GeV/c)}^{-2}$$

→ Interference visible as downturn for very low values of  $|t|$

See Spencer Klein's Talk

Ultra-peripheral collisions and hadronic structure,  
10 Feb 2017, 18:00



# Coherent Photoproduction in UPCs STAR

→ What does this look like in the UPC case?


→ Coherent production characterized by large slope parameter for small  $|t|$

$$\text{SLOPE} = 426 \pm 2 \text{ (GeV/c)}^{-2}$$


→ **Interference** visible as downturn for very low values of  $|t|$

See Spencer Klein's Talk


Ultra-peripheral collisions and hadronic structure,  
10 Feb 2017, 18:00



# Photon Interactions in 60-80% A+A?




# Photon Interactions in 60-80% A+A?





# Photon Interactions in 60-80% A+A?



# Novel Probe of the Medium? STAR

→ Vector meson ( $\rho^0$ ,  $J/\psi$ ) is coherently produced with very low  $p_T$

# Novel Probe of the Medium? STAR


- Vector meson ( $\rho^0$ ,  $J/\psi$ ) is coherently produced with very low  $p_T$
- Medium is produced by strong interaction

# Novel Probe of the Medium? STAR

- Vector meson ( $\rho^0$ ,  $J/\psi$ ) is coherently produced with very low  $p_T$
- Medium is produced by strong interaction
- Meson has very low  $p_T$  → it remains in the medium until it decays to an  $l^+ l^-$  pair

# Novel Probe of the Medium? STAR

- Vector meson ( $\rho^0$ ,  $J/\psi$ ) is coherently produced with very low  $p_T$
- Medium is produced by strong interaction
- Meson has very low  $p_T$  → it remains in the medium until it decays to an  $l^+l^-$  pair
- Since leptons are unaffected by the strong force, they carry information to the final state
- Provide a clean probe of medium



## Part I

- Very low  $p_T$   $e^+e^-$  invariant mass spectra in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV and U+U collisions at  $\sqrt{s_{NN}} = 193$  GeV


## Part II

- First look at  $\mu^+\mu^-$  invariant mass spectra with the Muon Telescope Detector at STAR in p+p collisions at  $\sqrt{s} = 200$  GeV and Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV



## Muon Telescope Detector (MTD)

- $|\eta| < 0.5$
- Azimuthal coverage  $\sim 45\%$
- Precise timing  $\sigma \sim 100$  picoseconds
- Precise spatial resolution  $\sim 1$  cm



# Motivation for STAR $\mu^+ \mu^-$



- STAR's Muon Telescope Detector allows for new studies of the dimuon continuum at RHIC energies with STAR.


Species	Energy (GeV)	~Sampled Luminosity
p+p (63% MTD)	500	28 pb <sup>-1</sup>
<b>p+p</b>	<b>200</b>	<b>122 pb<sup>-1</sup></b>
p+Au	200	409 nb <sup>-1</sup>
d+Au	200	94 nb <sup>-1</sup>
<b>Au+Au 2014</b>	<b>200</b>	<b>14 nb<sup>-1</sup></b>
Au+Au 2016	200	12 nb <sup>-1</sup>

## Why measure $\mu^+ \mu^-$ in central heavy ion collisions?



- Low mass excess is sensitive to the lifetime of the medium
- Rho meson broadening may be a probe of chiral symmetry restoration
- Intermediate mass region is sensitive to QGP thermal radiation.

Measurements of  $\mu^+ \mu^-$  in p+p collisions :

- Baseline for Au+Au measurements



# $\mu^+ \mu^-$ in Run14 Au+Au @ $\sqrt{s_{NN}} = 200$ GeV **STAR**



Data Triggered by dedicated Dimuon Trigger

In 60-80% Au+Au :

→ Clear  $\phi$  and  $J/\psi$  peaks

→  $S/B > \sim 1/10$

( $\sim 1/100$  to  $1/250$  in  $e^+e^-$ )

Significantly more data in semi-central and central collisions

→ **Expect exciting new  $\mu^+ \mu^-$  results from STAR**

## Low $p_T$ $e^+e^-$ measurements in Au+Au and U+U

- Significant excess yield with respect to hadronic sources observed across wide invariant mass range ( $M_{ee} < 4 \text{ GeV}/c^2$ )
- Shape and slope of  $t$  distribution show evidence of interference
- Large slope parameter is evidence for UPC-like coherent photoproduction mechanism
- Future studies will help clarify production mechanism
- May provide a novel probe of the medium


## First look at $\mu^+\mu^-$ invariant mass spectra from the Muon Telescope Detector at STAR

- New measurements possible at STAR with MTD
- Clear  $\omega$ ,  $\phi$ ,  $J/\psi$ ,  $\psi(2S)$  signal in p+p collisions @  $\sqrt{s} = 200 \text{ GeV}$
- Expect exciting new  $\mu^+\mu^-$  results from STAR

**Thank you**





$J/\psi$   $dN/dt$  distribution for Au+Au @  $\sqrt{s_{NN}} = 200$  GeV (40-80%)



See Wangmei Zha's poster : Poster ID 258

J/ $\psi$   $R_{AA}$  for Au+Au collisions @  $\sqrt{s_{NN}} = 200$  GeV and  
U+U collisions @  $\sqrt{s_{NN}} = 193$  GeV





PRL**116**, 222301 (2016)