

PREPARED FOR SUBMISSION TO JINST

20TH INTERNATIONAL CONFERENCE ON INSTRUMENTATION FOR COLLIDING BEAM PHYSICS

24 - 28 FEBRUARY, 2020

BUDKER INSTITUTE OF NUCLEAR PHYSICS, NOVOSIBIRSK, RUSSIA

The STAR Detector Upgrades for the BES-II and at Forward Rapidity

C. Yang¹ and Q. Yang on behalf of STAR Collaboration

*Key Laboratory of Particle Physics and Particle Irradiation (MOE),
Shandong University,
Qingdao 266237, China*

E-mail: chiyang@email.sdu.edu.cn

ABSTRACT: The Beam Energy Scan Phase II program (BES-II) at the Relativistic Heavy Ion Collider (RHIC) is examining the center-of-mass collision energy region from 7.7 GeV to 19.6 GeV which was determined from the results of BES-I. Key measurements such as the net proton kurtosis, the directed flow and the dilepton production are possible during BES-II with an order of magnitude better statistics due to the Low Energy RHIC electron Cooling (LEReC) upgrade of RHIC and the STAR detector upgrades. Beyond BES-II, the STAR Collaboration is currently designing, constructing, and installing a suite of new detectors in the forward rapidity region ($2.5 < \eta < 4$), enabling a program of novel measurements in pp, pA and AA collisions. To fully explore this physics, the forward upgrade needs superior detection capability for neutral pions, photons, electrons, jets and leading hadrons by adding charged particle tracking and electromagnetic and hadronic calorimetry to STAR's capabilities at forward rapidity. In this proceedings, we will present the details on the STAR detector upgrades for the BES-II and at forward rapidity. Physics in cold QCD program will also be discussed.

KEYWORDS: STAR detector upgrade, BES-II, Cold QCD, Forward rapidity upgrade, sTGC, HCal, ECal

¹Corresponding author.

Contents

1	STAR Detector Upgrades for RHIC Beam Energy Scan Phase II	1
2	Cold QCD Program	2
3	STAR Detector Upgrades at Forward Rapidity	2
3.1	Forward Tracking System	3
3.2	Forward Calorimeter System	3
4	Summary	4
5	Acknowledgement	5

1 STAR Detector Upgrades for RHIC Beam Energy Scan Phase II

The Relativistic Heavy Ion Collider (RHIC) at BNL is uniquely suited to study the phase diagram of strongly interacting nuclear matter, search for the critical point and find evidence of a first-order phase transition through its Beam Energy Scan (BES) program. The BES-I program has localized the most interesting regions (7.7 GeV - 19.6 GeV) and has identified the observables which are likely to be the most discriminating for understanding the QCD phase structure. However, several of the key measurements such as the net proton kurtosis, the directed flow and the dilepton production were found to require higher statistics in order to provide a quantitative physics conclusion [1]. Therefore, the second phase of the beam energy scan (BES-II) started from 2019 with the upgraded Low Energy RHIC electron Cooling (LReC) and various sub-system of the STAR detector. The STAR detector upgrades for BES-II are consist of inner TPC (iTPC), Event Plane Detector (EPD), and the endcap TOF (eTOF) upgrades [2], as shown in FIG. 1.

The STAR iTPC upgrade is to replace all the inner-sector wires which are aging with continuous pad rows [3]. The upgrade will provide better momentum resolution, lower p_T threshold from 125 MeV/c to 60 MeV/c, better dE/dx resolution, improved acceptance at high rapidity from 1.0 to 1.5 and higher track reconstruction efficiency [1]. The EPD serves to improve event-plane reconstruction for heavy-ion flow measurements, the collision centrality and trigger detection for events. A new EPD can cover the pseudo-rapidity range between 2.1 and 5.1 and replace Beam Beam Counter(BBC) [1, 4]. The eTOF upgrade will extend particle identification (PID) capability and cover the pseudorapidity region of $2.5 < \eta < 4$. The fixed-target program at STAR with the eTOF upgrade will enable the energy scan to extend below 7.7 GeV [5]. All of these three sub-detectors are fully operational in STAR BES-II program.




Figure 1. The layout of STAR detector upgrades for RHIC BES-II program.

2 Cold QCD Program

The detector upgrades for BES-II program have substantially enhance STAR's already excellent capabilities. These upgrades will enable STAR to continue its unique, ground-breaking mid-rapidity science program in the period following BES-II. The key physics opportunities envisioned address three broad areas of interest within the cold QCD community in the years following the BES-II. These programs will shed light on the dynamics of low and high x partons in cold nuclear matter (CNM) and how the fragmentation and hadronization of these partons is modified through interactions within the CNM and experiments to study the 2+1d spatial and momentum structure of protons and nuclei. These measurements will provide critical new insights into the QCD structure of nucleons and nuclei in the near term, as well as the high precision data that will be essential to enable rigorous universality tests when combined with future results from the Electron Ion Collider.

The transverse polarized proton colliding with unpolarized proton and the longitudinal polarized proton-proton collision at $\sqrt{s}=510$ GeV are scheduled in 2020 and 2021 at RHIC and STAR will be the only operational detector at RHIC during this period. The scientific goals are study the parton transverse momentum distribution (TMDs) at low and high x as well as gluon distribution at small x . With a delivered luminosity of 1.1 fb^{-1} , it will give unprecedented precision for several polarized observables such as transverse spin asymmetry (A_{UT}) for Collins observables [6].

The transverse polarized proton-nuclear collisions at $\sqrt{s}=200$ GeV are scheduled in parallel with sPHENIX running, which including polarized proton with unpolarized proton, polarized proton-Au, and polarized proton-Al collisions. These collisions system are an ideal system to test gluon hadronization effects in cold nuclear matter at high collision energy and Q^2 accessed at RHIC which will provide a clearer picture of modified hadronization. The high statistics polarized p+Au dataset and A scan can precisely determine the mass dependence of nuclear hadronization effect [6].

3 STAR Detector Upgrades at Forward Rapidity

The STAR detector upgrades at forward rapidity is ongoing to support the cold QCD program, together with other interesting topics in forward rapidity. The upgrades are consist of Forward

Calorimeter System (FCS) and Forward Tracking System (FTS) [7].

3.1 Forward Tracking System

FTS includes 3 silicon disks and 4 small-strip Thin Gap Chamber (sTGC) disks, providing 7 tracking points in η coverage from 2.5 to 4.

The silicon tracker with fine granularity in ϕ and coarse granularity in r is using the single-sided double metal AC-coupled architecture. Each silicon disk has 12 modules. The silicon sensors have good charge separation, momentum resolution and precise position resolution to meet the requirement of the FTS [9]. It will be placed at STAR center $z=139.9\text{cm}$, limited by the requirement of η coverage from 2.5 to 4. The sensor signal is read out at its edge. The frontend readout chips are APV25-S1 in FTS [9]. Each APV25-S1 has 128 channels, which have the function of pre-amplifier and shaper. 288 APV chips will be used in 3 silicon disks.

The sTGC is a gaseous detector based on the technology developed by the ATLAS collaboration [8]. The position resolution is designed at the level of $\sim 100\mu\text{m}$. The STAR sTGC project is planned to build 4 layers of sTGC, 4 modules per layer. Each sTGC module consists of two chambers which are perpendicular to each other for the position measurements in x-y coordinate. First $30\text{cm}\times 30\text{cm}$ size pre-prototype was built at Shandong University and installed at STAR in June 2019. As shown in FIG. 2, simply taking the ratio of the signal counts with the coincidences between different chambers, the efficiency achieved in test beam is $>98\%$. A $60\text{cm}\times 60\text{cm}$ full size prototype has been built in late 2019 and the signal coincidences shown in FIG. 2 are consistent with that in pre-prototype. Mass production is planned to start in 2020 and all 20 modules of sTGC (16 + 4 spares) will be finished in year 2021.




Figure 2. The signal coincidences of different sTGC chambers for efficiency check. Left: Pre-prototype in STAR 2019 data-taking. Right: Full size prototype in cosmic ray test.

3.2 Forward Calorimeter System

The FCS consisted of Electromagnetic Calorimeter (ECal) and Hadronic Calorimeter (HCal). The ECal reuses the lead-scintillator calorimeter from the PHENIX Collaboration with new SiPM based readout. Penetrating wavelength-shifting fibers for light collection. The HCal is the first hadronic calorimeter in STAR which is an ion-scintillator sandwich sampling calorimeter with the same

SiPM readout as ECal. Intensive test beam for Forward Calorimeters at FNAL has been done in 2019. As shown in FIG. 3, performances of ECal and HCal prototype are close to the requirements.




Figure 3. The first prototype including the HCal, the ECal, pre-shower, and the sTGC was installed for the 2019 data-taking in STAR.




Figure 4. (a) ECal prototype energy resolution. (b) HCal prototype energy resolution.

4 Summary

The STAR detector upgrades, including the iTPC, the EPD, and the eTOF upgrades, have been finished and all three sub-detectors are fully operational for RHIC BES-II program. A forward rapidity upgrade which is consist of tracking detectors (silicon and sTGC) and calorimeters (ECal and HCal) at STAR is ongoing to further investigate the nature of QCD, especially the cold QCD physics at both very high and very low partonic momentum fraction. Current prototype test shows that the prototype performances meet the STAR forward rapidity upgrade requirements.

5 Acknowledgement

The work is partly supported by the National Natural Science Foundation of China (No. 11890713), the Shandong Provincial Natural Science Foundation, China (No. ZR2019MA005) and the Program of Qilu Young Scholars of Shandong University.

References

- [1] STAR Collaboration, *Studying the Phase Diagram of QCD Matter at RHIC*, STAR Note 0598, <https://drupal.star.bnl.gov/STAR/starnotes/public/sn0598>
- [2] C. Yang, on behalf of STAR Collaboration, *The STAR Beam Energy Scan Phase II Physics and Upgrades*, *Nucl.Phys.A* **967** (2017) 800-803
- [3] F. Shen, S. Wang, C. Yang and Q. Xu, on behalf of STAR Collaboration, *MWPC Prototyping and Testing for STAR inner TPC Upgrade*, *JINST* **12** (2017) no.06, C06008
- [4] STAR Collaboration, *An Event Plane Detector for STAR*, STAR Note 0666, <https://drupal.star.bnl.gov/STAR/starnotes/public/sn0666>
- [5] STAR Collaboration and CBM eTOF Group, *Physics Program for the STAR/CBM eTOF Upgrade* arXiv:1609.05102 [nucl-ex]
- [6] E. Aschenauer, et al., *The RHIC Cold QCD Plan for 2017 to 2023: A Portal to the EIC*, arXiv:1602.03922 [nucl-ex]
- [7] Q. Yang, on behalf of STAR Collaboration, *The STAR BES-II and Forward Rapidity Physics and Upgrades*, *Nucl.Phys.A* **982** (2019) 951-954
- [8] A. Abusleme, et al., *Performance of a Full-Size Small-Strip Thin Gap Chamber Prototype for the ATLAS New Small Wheel Muon Upgrade*, *Nucl.Instrum.Meth.A* **817** (2016) 85-92
- [9] STAR Collaboration, *The STAR Forward Calorimeter System and Forward Tracking System*, STAR Note 0648, <https://drupal.star.bnl.gov/STAR/starnotes/public/sn0648>