


# Hypernuclei Production in Heavy-Ion Collisions


(at finite baryon density)



**CPOD 2024**

**Yue Hang Leung**  
for the STAR collaboration

University of Heidelberg  
20<sup>th</sup> May, 2023



## Outline

- Introduction
- ${}^3_{\Lambda}\text{H}$  Yields and Particle Ratios
- Other Observables
  - ${}^4_{\Lambda}\text{H}$  Yields
  - Collective Flow
- Summary
- Outlook



# What can hypernuclei production in heavy-ion collisions tell us about the QCD phase diagram?

- Hypernuclei yields have been suggested to be sensitive to the **onset of deconfinement**


- $S_3 = \frac{{}^3_{\Lambda}\text{H}}{{}^3\text{He} \times \frac{\Lambda}{p}}$  may be enhanced in

systems involving partonic interactions

S. Zhang et al, Phys. Lett. B 684 (2010) 224


- Baryon clustering near critical point may lead to enhancement of light nuclei ( $A \geq 3$ ) yields

E. Shuryak et al, Phys. Rev. C 101 (2020) 034914




# What is the role of hyperon-nucleon (YN) interaction in the equation-of-state of high baryon density matter?

- **Hyperon Puzzle:** difficulty to reconcile the measured masses of neutron stars with the presence of hyperons in their interiors



?




*Can hypernuclei production be used to constrain the in-medium Y-N interaction?*

- Density dependent YN, YNN interactions are essential for solving the hyperon puzzle

How and when are light nuclei formed in heavy ion collisions?


Need a solid understanding in  
**hypernuclei production mechanisms**  
before we can use them as **probes for**  
**medium properties**

# What Have We Learnt From Light Nuclei Production?



STAR, Phys. Rev. Lett. 130 (2023) 202301

- d/p is fairly well described by thermal model, but t/p is overestimated





ALICE, Phys. Rev. C 107 (2023) 064904

ALICE, arXiv:2311.11758v1

- $^4\text{He}/p$  is well described by thermal model at LHC energies, but underestimated by various implementations of coalescence formation

**Recent data poses challenges for nuclei production models**

# Hypertriton ( ${}^3_{\Lambda}\text{H}$ ) and Hyperhydrogen-4 ( ${}^4_{\Lambda}\text{H}$ )



**$\Lambda$  binding energy**

~0.1 MeV

~2.2 MeV

**Excited states**


Not observed




- Due to its very small binding energy,  ${}^3_{\Lambda}\text{H}$  production provides unique input for nuclei production models

# STAR Beam Energy Scan II




- BES-I (2009-2011)
  - Au+Au collisions  $\sqrt{s_{NN}} = 7.7-62$  GeV
  - Main objectives:
    - **Search for onset of deconfinement**
    - **Search for critical end point**



- BES-II (2017-2021)
  - High statistics Au+Au collisions  $\sqrt{s_{NN}} = 3-54.4$  GeV
  - Fixed target (FXT) collisions extend energy reach down to  $\sqrt{s_{NN}} = 3$  GeV
    - **Search for possible formation and investigate properties of dense baryonic matter**




# ${}^3_{\Lambda}\text{H}$ Rapidity and $p_T$ Spectra



- Measurements cover **11 different energies** from  $\sqrt{s_{NN}} = 3$  to 27 GeV


# ${}^3_{\Lambda}\text{H}$ Excitation Function



- Steep increase from  $\sqrt{s_{\text{NN}}} = 27$  to 4 GeV
- Plateaus at  $\sqrt{s_{\text{NN}}} = 3-4$  GeV
- Interplay between increasing baryon production and stronger strangeness canonical suppression towards low energies

**Establishes low energy collision experiments as a promising tool to study exotic strange matter**


# Nuclei-to-Hadron Ratios



- ${}^3_{\Lambda}H/\Lambda$  ratio in a thermal model calculation is independent of volume and strangeness correlation length
- ${}^3_{\Lambda}H/\Lambda$ , similar to  $t/p$ , are overestimated by thermal model by a factor of 2

**${}^3_{\Lambda}H$  (and  $t$ ) yields are not in equilibrium and fixed at chemical freeze-out along with other light hadrons**

# Mean Transverse Momentum




- ${}^3_{\Lambda}$ H and  $t$  have similar mean  $p_T$
- Both  ${}^3_{\Lambda}$ H and  $t$  tend to have lower mean  $p_T$  than the blast-wave parametrization using measured kinetic freeze-out parameters from light hadrons ( $\pi, K, p$ )

**${}^3_{\Lambda}$ H (and  $t$ ) do not follow same collective expansion as light hadrons**

- Softening of the  ${}^3_{\Lambda}$ H may be explained by coalescence model with Wigner-function formalism

D. Liu et al, arXiv:2404.02701

# Mean Transverse Momentum




- The mean  $p_T$  for  $\sqrt{s_{NN}} = 3-4.5$  GeV and  $\sqrt{s_{NN}} = 7.7-27$  GeV seem to exhibit two different trends

Change in medium properties or expansion dynamics?


see Y. Zhou, 17:00 20/05 (Mon.)

# Centrality Dependence of ${}^3_{\Lambda}\text{H}$ Production




- The yield in mid-central (10-40%) collisions follows the same trend as central (0-10%) collisions

# Centrality Dependence of ${}^3_{\Lambda}\text{H}$ Production



- ${}^3_{\Lambda}\text{H}$  production increases more steeply compared to  $N_{\text{part}}$ , seems to be more apparent below 7.7 GeV

# Centrality Dependence of ${}^3_{\Lambda}\text{H}$ Production





- Proton yield scales with  $N_{\text{part}}$
- $\Lambda$  yield increases more steeply than  $N_{\text{part}}$ , particularly at low collision energies

see Y. Zhou, 17:00 20/05 (Mon.)

- At low energies,  ${}^3_{\Lambda}\text{H}$  production tends to increase more steeply than proton,  $\Lambda$ ,  ${}^3\text{He}$

**Suppression of  ${}^3_{\Lambda}\text{H}$  production in more peripheral collisions at low energies**


# Centrality Dependence of ${}^3_{\Lambda}\text{H}$ Production



- Similar observation in Ag+Ag collisions at  $\sqrt{s_{NN}}=2.55$  GeV

**Suppression related to the nature of the created medium?**


# Strangeness Population Factor $S_3$



- An enhancement of  $S_3$  was proposed as a probe for deconfinement


S. Zhang et al, Phys. Lett. B 684 (2010) 224

# Strangeness Population Factor $S_3$



- An enhancement of  $S_3$  was proposed as a probe for deconfinement
- Data indicates a mild increase in  $S_3$ , do not follow the expectations of the model


# Strangeness Population Factor $S_3$



- The measured  $S_3$  is close to thermal model predictions
- The increasing trend is driven by the decreasing feed-down from  ${}^3\text{He}$  towards higher energies

T. Reichert et al, Phys.Rev.C 107(2023)014912

# Strangeness Population Factor $S_3$





- UrQMD + Coalescence seem to overshoot the data
- A key prediction from coalescence models is the suppression of  ${}^3_{\Lambda}H$  production in small systems due to its large radius

T. Reichert et al, Phys.Rev.C 107(2023)014912

- Best represented by investigating the multiplicity dependence, since  $dN_{ch}/d\eta$  is a good proxy for volume
- Possible feed-down should be accounted for when interpreting results


E864, Phys. Rev. C 70(2004)024902  
 STAR, Science 328(2010)58  
 STAR, arXiv:2310.12674  
 ALICE, Phys. Lett. B 754(2016)360

# Multiplicity Dependence of $S_3$ (stable nuclei)



- Unstable nuclei production are suppressed relative to stable nuclei (see backup)
- The true value of  $S_3$  (stable nuclei) very likely lies between the upper and lower limits

# Multiplicity Dependence of $S_3$ (stable nuclei)




$$S_3 \text{ (stable nuclei)} \approx 0.35$$

- Existing data for  $S_3$  considering stable nuclei only do not exhibit significant dependence on collision energy, system size
- Data show **milder multiplicity dependence** compared to coalescence, particularly 3-body
- Thermal model tends to overpredict  $S_3$  at  $dN_{ch}/d\eta=200$  or lower

*More data at very low and very high  $dN_{ch}/d\eta$  is needed*

# ${}^4_{\Lambda}\text{H}$ Production




- Non-monotonic behavior of hypernuclei to nuclei yields vs mass number

**Suggestive of creation of unstable hypernuclei**




# ${}^4_{\Lambda}\text{H}$ Production



- ${}^4_{\Lambda}\text{H}$  yields tend to be closer to thermal model predictions compared to  ${}^3_{\Lambda}\text{H}$

Size of hypernuclei plays a role in their production?

# Hypernuclei Collective Flow



STAR, Phys.Lett.B 827(2022)136941  
STAR, Phys. Rev. Lett. 130(2023)212301


- Directed flow of hypernuclei follows mass scaling
- JAM + coalescence approx. describes the data

**Qualitatively consistent with coalescence formation of hypernuclei**

# Summary

- ${}^3_{\Lambda}\text{H}$  yields in central collisions overestimated by thermal model by a factor of 2
- ${}^3_{\Lambda}\text{H}$  mean  $p_T$  tends to be lower than blast-wave parametrization from light hadrons
  - ${}^3_{\Lambda}\text{H}$  is not in thermal equilibrium / does not decouple at same time as light hadrons
- Data for  $S_3$  (stable nuclei) are consistent with flat or slightly increasing trend with  $dN_{\text{ch}}/d\eta$ 
  - Milder multiplicity dependence compared to coalescence models
- Suppression of  ${}^3_{\Lambda}\text{H}$  in 10-40% collisions at low collision energies observed
- ${}^4_{\Lambda}\text{H}$  yields tend to be closer to thermal model predictions than  ${}^3_{\Lambda}\text{H}$ 
  - Data provide new input to investigate system size and nuclei size dependence of nuclei production
- ${}^3_{\Lambda}\text{H}$  mean  $p_T$  seem to exhibit two separate trends for  $\sqrt{s_{NN}} = 3 - 4.5\text{GeV}$  and  $7.7 - 27\text{GeV}$ 
  - Change in medium properties or expansion dynamics?

# Outlook



## RHIC-STAR

- Heavier hypernuclei, including  ${}^4_{\Lambda}H$ ,  ${}^4_{\Lambda}He$ ,  ${}^5_{\Lambda}He$ ,  ${}^6_{\Lambda}H$  at FXT energies
- High statistics data at RHIC top energy give opportunities for multiplicity dependence study



## FAIR-CBM and HIAF

- Double- $\Lambda$  hypernuclei to constrain  $\Lambda$ - $\Lambda$  interaction, essential for hyperon puzzle resolution

Thank you for listening!



# Feed-down from Unstable Nuclei




- Feed-down fractions estimated using Thermal-FIST

V. Vovchenko et al, Phys. Lett. B 809 (2020) 135746

- Feed-down correction of  $S_3$  from unstable nuclei estimated using Thermal-FIST


# Feed-down from Unstable Nuclei



Phys. Rev. C 65 (2001) 014906

- Suppression of  $A=4$  unstable states compared to  ${}^4\text{He}$  ground state observed at E864

# Centrality Dependence of ${}^4_{\Lambda}$ H Production



- Within uncertainties, the centrality dependence of  ${}^4_{\Lambda}$ H production is consistent to that of  ${}^3_{\Lambda}$ H at mid-rapidity