MULTI-DIMENSIONAL MEASUREMENTS OF THE PARTON SHOWER IN PP COLLISIONS AT RHIC ABSTRACT FOR QM 22 POSTER

MONIKA ROBOTKOVA
(FOR THE STAR COLLABORATION)
NUCLEAR PHYSICS INSTITUTE, CZECH ACADEMY OF SCIENCES

Jets are collimated sprays of hadrons and serve as an experimental tool for study-7 ing the dynamics of quarks and gluons. In particular, differential measurements of 8 jet substructure enable a systematic exploration of the parton shower evolution. 9 The SoftDrop grooming technique utilizes the angular ordered Cambridge/Aachen 10 reclustering tree and provides a correspondence between the experimental observ-11 ables such as the shared momentum fraction (z_g) , groomed jet radius or split 12 opening angle (R_g) and the QCD splitting functions in vacuum. In this poster, 13 we present fully corrected correlations between $z_{\rm g}$ and $R_{\rm g}$ at the first split for 14 jets of varying momenta and radii in pp collisions at $\sqrt{s} = 200$ GeV. To study 15 the evolution along the jet shower, we also present the splitting observables at 16 the first, second and third splits along the jet shower for various jet and initiator 17 prong momenta. As these novel measurements are presented in three dimensions, 18 we outline the correction procedure so that it can be used as a template for future 19 multi-differential measurements across all experiments. 20

Date: November 22, 2021.