



Elliptic flow of light nuclei in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV

Chitrasen Jena

for the STAR collaboration

Institute of Physics, Bhubaneswar



High Energy Physics Symposium 2010

December 13-18, 2010, Jaipur, India



December 13-18,
2010


XIX DAE-BRNS HEP Symposium,
Jaipur, India

Outline

- Introduction
- Experiment
- Analysis Method
- Results and Discussions
- Summary



Introduction



$$\varepsilon = \frac{\langle y^2 - x^2 \rangle}{\langle y^2 + x^2 \rangle}$$

Coordinate-Space
Anisotropy



Momentum-Space
Anisotropy

$$E \frac{d^3 N}{d^3 p} = \frac{1}{2\pi} \frac{d^2 N}{p_T dp_T dy} \left(1 + 2 \sum_{n=1}^{\infty} v_n(p_T, y) \cos(n(\phi - \Psi_r)) \right)$$

$$v_n = \langle \cos(n(\phi - \Psi_r)) \rangle \quad \phi = \tan^{-1} \left(\frac{p_y}{p_x} \right)$$

➤ Elliptic flow (v_2) is a good probe of the early stage of the collision.




Motivation

- Light nuclei and anti-nuclei are formed through coalescence of nucleons (produced/participant) and anti-nucleons^{*}. This formation process is generally believed to happen at a late stage of the evolution due to their small binding energy.
- Production of light nuclei provides information of space-momentum correlation among these nucleons.
- By studying the v_2 of light nuclei and comparing to that of their constituent nucleons, we will have a better understanding of coalescence process for hadronization.

^{*}H.H. Gutbrod et al., Phys. Rev. Lett. 37, 667 (1976).



STAR Detector



Full azimuthal particle identification!




December 13-18,
2010

XIX DAE-BRNS HEP Symposium,
Jaipur, India

Data Set

Data : Run 7 Au+Au 200 GeV
Number of events: ~62 M
(~2.5 times more statistics
compared to previous run)

➤ Measurement of the ionization energy loss (dE/dx) of charged tracks in the TPC gas are used to identify the light nuclei.





Analysis Method

➤ ^3He signal is almost background free for $p_T > 1.4 \text{ GeV}/c$.

$$Z = \ln\left(\frac{(dE/dx)_{\text{measure}}}{(dE/dx)_{\text{predict}}}\right)$$

➤ TPC is used to determine the event plane.





$$\frac{dN}{d\phi} \propto 1 + 2v_2^{obs} \cos[2(\phi - \Psi_R)]$$

$$v_2^{final} = \frac{v_2^{obs}}{R_2}$$

R_2 is the second order event plane resolution



v_2 vs. p_T





STAR, arXiv:0909.0566 [nucl-ex]

- v_2 of ${}^3\text{He}$ is well described by the dynamical coalescence model.
- Anti-triton v_2 has been shown for $0.3 < p_T < 1.2$ GeV/c.

Model Calculations: S. Zhang et al., Phys. Lett. B 684 (2010) 224



NCQ Scaling





STAR, PRC 72, 014904 (2005)
 STAR, arXiv:0909.0566 [nucl-ex]
 PHENIX, PRL 99, 052301 (2007)

$$d(p+n) : n_q = 2 \times 3 \quad {}^3\text{He}(2p+n) : n_q = 3 \times 3$$

- v_2 of ${}^3\text{He}$ seem to follow the atomic mass number (A) scaling.
- Number of constituent quark (NCQ) scaling holds good for v_2 of ${}^3\text{He}$.



Centrality Dependence



Similar to other hadrons, at more central collision the larger value of v_2/ϵ indicates stronger collective expansion.

STAR, PRC 77, 054901 (2008)




December 13-18,
2010

XIX DAE-BRNS HEP Symposium,
Jaipur, India

10

Mass Dependence



- Both v_2 and $\langle p_T \rangle$ trends are consistent with expectations from Blast-wave (BW) model fit.
- v_2 values up to ${}^3\text{He}$ mass has reasonable agreement within the Blast-wave formalism but differences seen in $\langle p_T \rangle$ beyond proton mass.

➤ *Statistical and systematic uncertainties are under study.*

π, p spectra: STAR, PRL 97, 152301 (2006)

$\pi, p v_2$: STAR, PRC 72, 014904 (2005)
STAR, PRC 77, 054901 (2008)

ϕ spectrum and v_2 : STAR, PRL 99, 112301 (2007)


${}^3\text{He}$ spectrum: STAR, arXiv:0909.0566 [nucl-ex]

BW: Z. Tang et al., PRC 79, 051901 (2009)
M. Shao et al., JPG 37, 085104 (2010)



Summary

- v_2 of light nuclei are measured in Au+Au@200 GeV using event plane method.
- p_T dependence of v_2 is well described by Dynamical Coalescence Model.
- v_2 of light nuclei seem to follow the atomic mass number scaling.
- Number of constituent quark scaling holds good for v_2 of light nuclei.
- At more central collision the large value of v_2/ϵ indicates stronger collective expansion.
- Both v_2 and $\langle p_T \rangle$ trends are consistent with expectations from Blast-wave model fit.



Backup



December 13-18,
2010

XIX DAE-BRNS HEP Symposium,
Jaipur, India

13

v_2 Comparison

