



# Reconstruction of $K^*(892)$ Resonance in Au+Au Collisions at 200 GeV at STAR

He Zheng\* (UCLA)  
for the STAR Collaboration

DNP2016

October 13-16, 2016  
Sheraton Vancouver Wall Centre  
Vancouver, BC Canada

The Relativistic Heavy Ion Collider (RHIC) produces a hot, dense and de-confined Quantum Chromodynamics (QCD) medium, called the quark-gluon plasma (QGP), with Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. The  $K^*(892)$  resonance is a short-lived vector meson with a life-time of 4 fm/c, shorter than the expected life-time of the QGP. The decay of the  $K^*$  and its properties may provide an effective tool to probe the evolution of the QGP produced. Experimentally,  $K^*$  is not a well-studied particle at STAR previously because of its fast decay and large combinatorial background. In recent years, improvements in data sample statistics and particle identification capability promise better  $K^*$  measurements. In this presentation, we report the reconstruction of invariant mass of  $K^*$  resonance via the hadronic decay channel  $K^*(892) \rightarrow K_S^0 \pi^\pm$  as a function of transverse momentum ( $p_T$ ) up to 5 GeV/c for various collision centrality classes. Physics implications of our measurements will also be discussed.

## Introduction

$K^*(892)$  candidate is reconstructed by inverting decay mode to obtain the distribution of invariant mass of the decay parent.


By special relativity,

$$m_{K^*} = \sqrt{E_{K^*}^2 - \vec{p}_{K^*}^2} = \sqrt{(E_{K_S} + E_\pi)^2 - (\vec{p}_{K_S} + \vec{p}_\pi)^2} \quad (c = 1)$$

So we should expect to observe a signal around 0.892 GeV/c<sup>2</sup>.

## Background Method:



Mixed-Event Background – Build reference background distribution by pairing decay daughters from different collision events to eliminate possible correlation dependence.



- The data used in this analysis were minimum bias trigger Au+Au collisions at 200 GeV collected in the Run 2011 from the STAR experiment.
- Particle Identification: TPC (Time Projection Chamber) dE/dx and TOF (Time of Flight) are used for pion identification.

## $K_S^0$ signals

Observed in the  $\pi^+\pi^-$  invariant mass distribution reconstructed from the decay topology method.



PDG value:  $m = 497.614 \pm 0.024$  MeV/c<sup>2</sup>


- $K^*(892)$  signals fitting:** The Breit-Wigner function with background fitted by a 3<sup>rd</sup> order polynomial.

$$\frac{Y * 0.01 * W}{2\pi((x - M)^2 + W^2/4)} + ax^3 + bx^2 + cx + d$$


where Y = yield, W = width, M = mass.

Examples of foreground (red) and event mixing background (blue):

Centrality 70%~80%,  $p_T = 4-5$  GeV/c




Centrality 60%~70%,  $p_T = 4-5$  GeV/c




## Results

- $K^*(892)$  signals:** Mixed-event background has been subtracted.


$K^*$  signals for  $p_T = 0.5-3$  GeV/c, all centrality combined




$p_T = 0.5-1$  GeV/c, centrality 50%~80%




$p_T = 0.5-1$  GeV/c, centrality 20%~50%




$p_T = 1-2$  GeV/c, centrality 50%~80%




$p_T = 1-2$  GeV/c, centrality 20%~50%




$p_T = 2-5$  GeV/c, centrality 50%~80%




$p_T = 2-5$  GeV/c, centrality 20%~50%



$K^*$  invariant mass vs  $p_T$



$K^*$  width vs  $p_T$



Possible sources of the differences from the PDG values may include the uncorrected efficiency and the effect of strong magnetic field created.

## Summary and Outlook

- The signals for  $K^*(892)$  resonance produced in Au+Au collisions at 200 GeV at STAR are significant. The data analysis confirms the existence of a measurable amount of  $K^*$ , which allows further study of its properties.
- Future study of new physics if possible, includes resonance decays in strong magnetic field. For example, how  $K^*$  mass changes with the magnetic field.

## Acknowledgement

Thanks to Prof. Huan Z. Huang for mentorship and to Dr. Gang Wang and the STAR collaboration for guidance. Thanks to Roli Esha, Liwen Wen for valuable help on my coding.

## Reference

[1]. STAR Collaboration, arXiv:nucl-ex/0412019v2, 22 Apr 2005

\*e-mail: peterzheng@ucla.edu