

APS April Meeting

January 28-31, 2017

Washington, DC



U.S. DEPARTMENT OF
ENERGY

DOE NP contract: DE-SC0013405

Measurements of single-spin asymmetries, A_L for W^\pm boson production in longitudinally polarized proton-proton collisions at STAR

• Motivation


• Theoretical Aspect

• Experimental Aspect


• Analysis

• Results


• Summary



Devika Gunarathne
(for the STAR Collaboration)
Temple University




MOTIVATION : Proton Helicity Structure



1989 : EMC : DIS

$$\Delta\Sigma = 0.12 \pm 0.09 \pm 0.14$$

“Spin Crisis”



Naive Parton Model

$$\frac{1}{2} = \frac{1}{2}(\Delta u_v + \Delta d_v)$$


Gluons , Sea quarks are polarized.
 $\xrightarrow{\text{Parton orbital angular momentum.}}$

Current Understanding

$$\langle S_z \rangle = \frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L$$

$$\Delta\Sigma = \int (\Delta u + \Delta d + \Delta s + \Delta\bar{u} + \Delta\bar{d} + \Delta\bar{s})dx$$

MOTIVATION : Current Knowledge of PDFs





$$\Delta\Sigma = \int (\Delta u + \Delta d + \Delta s + \Delta\bar{u} + \Delta\bar{d} + \Delta\bar{s})dx$$

DIS


- Well measured!
- Not sensitive to flavor separation!

SIDIS


- FF's use to tag flavor!
- Flavor separation / quark, anti-quark separation!
- But large uncertainties in FFs.



Large uncertainties w.r.t quark PDFs




Phys. Rev. D 82, 114018 (2010)



Theoretical Aspects - $W A_L$

- Probing quark / anti-quark (sea) flavor structure using W boson production at RHIC

W production in p+p,



$$A_L = \frac{\sigma^+ - \sigma^-}{\sigma^+ + \sigma^-}$$


$$A_L^{e^-} \approx \frac{\int_{\otimes(x_1, x_2)} [\Delta \bar{u}(x_1) d(x_2) (1 - \cos \theta)^2 - \Delta d(x_1) \bar{u}(x_2) (1 + \cos \theta)^2]}{\int_{\otimes(x_1, x_2)} [\bar{u}(x_1) d(x_2) (1 - \cos \theta)^2 + d(x_1) \bar{u}(x_2) (1 + \cos \theta)^2]}$$

$$\langle x_{1,2} \rangle \sim \frac{M_W}{\sqrt{s}} e^{\pm \eta_e/2}$$

$$\eta = -\ln \left(\tan \left(\frac{\theta}{2} \right) \right)$$

In comparison to SIDIS,

- Direct sensitivity to \bar{u} , \bar{d} .
- Large Q^2 defined by W mass (more reliable perturbative calculation / higher twist effects unimportant!).
- Parity violating coupling gives rise to single-spin asymmetry which is directly related to anti-quark helicity PDFs.
- Free of FFs.
- Easy detection via decay leptons.




EXPERIMENTAL ASPECT - RHIC

- RHIC : Relativistic Heavy Ion Collider**


The World's first polarized hadron collider!

Spin varies from bunch to bunch. Spin pattern changes from fill to fill. Spin rotators provide choice of spin orientation.



EXPERIMENTAL ASPECT - STAR


• STAR : Solenoidal Tracker At RHIC



TPC : $-1.3 < \eta < +1.3$
 BEMC : $-1.0 < \eta < +1.0$
 EEMC : $+1.1 < \eta < +2.0$
 FGT : $+1.0 < \eta < +2.0$


TPC: **Charged particle tracking**
 BEMC, EEMC: **EM Calorimetry**

RHIC p+p runs : Luminosity





Run	L (pb^{-1})	P (%)	FOM (P^2L) (pb^{-1})
2009	12	0.38	1.7
2011	9.4	0.49	2.3
2012	77	0.56	24
2013	246.2	0.56	77.2


ANALYSIS -Mid rapidity STAR W selection criteria




- Isolated high P_T track pointing to isolated EMC cluster.
- Large Imbalance in the reconstructed vector P_T sum in 4π due to undetected neutron.



TPC track extrapolated
to Barrel colorimeter tower grid



Transverse plane view



- Several tracks pointing to several EMC clusters.
- Vector P_T sum is balanced by the Jet opposite in π .

• Mid-rapidity STAR W selection criteria


- Match $P_T > 10$ GeV track to BEMC cluster
- Isolation ratio 1 / Isolation ratio 2
- P_T -balance cut

$$E_T^e / E_T^{4X4} > 95\%$$


$$E_T^e / E_T^{\Delta R < 0.7} > 88\%$$

$$\vec{p}_T^{bal} = \vec{p}_T^e + \sum_{\Delta R > 0.7} \vec{p}_T^{jets}$$

$$P_T\text{-balance } \cos(\phi) = \frac{\vec{p}_T^e \cdot \vec{p}_T^{bal}}{|\vec{p}_T^e|}$$



ANALYSIS -Mid rapidity STAR W BG Estimation



Primary Background

- Data-driven QCD : BG Events which satisfy $e^{+/-}$ candidate isolation cuts due to “jet” escape detection outside STAR acceptance , $|\eta| > 2$.
- Second EEMC : due to “jet” escape detection at “non-existent” East EEMC, estimate based on “real” West EEMC

ElectroWeak Background



- Determine from MC simulation

$$Z \longrightarrow e^+ + e^-$$

$$W \longrightarrow \tau + \nu$$

RESULTS - W A_L - STAR 2011+2012 (published)

- STAR 2011 + 2012 W A_L Published Results




Impact on PDFs : DSSV

Only DIS + SIDIS data

E. Aschenauer, et.al. arXiv:1304.0079


DIS + SIDIS + STAR W A_L data



- A_L for W^+ is consistent with theoretical predictions constrained by polarized SIDIS data.
- A_L for W^- is larger than the prediction for $\eta_e < 0$, which suggest large $\Delta\bar{u}$.
- Indication of positive $\Delta\bar{u}$ at $0.05 < x < 0.2$.

RESULTS - W A_L - STAR 2011+2012 vs 2013

- STAR 2013 W A_L Preliminary Results in comparison to STAR 2011+2012 published results



- STAR 2013 W A_L Preliminary results is the **Most Precise** measurements of W A_L up to date!
- STAR 2013 preliminary W A_L results **consist** with published 2011 + 2012 results.
- Uncertainties were **reduced by 40 %**.

SUMMARY

- W boson production in longitudinally polarized p+p collisions at RHIC is a unique tool to probe quark antiquark helicity PDFs of the nucleon.
- Mid-rapidity (Run 11/12): Published W longitudinal single spin asymmetry results suggest large anti-u quark polarization along with broken QCD sea.
- New prelim. result of STAR 2013 $W A_L$ is the most precious measurement up to date. These results will help to further constrain antiquark helicity distributions.
- New STAR 2013 $W A_L$ prelim. results consistent with published STAR 2011+2012 results.